국내 육성 벼 243 품종의 형태적 특성에 따른 품종 분류

김미선¹ · 이해정^{1,2} · 유달아¹ · 송재영¹ · Marjohn Nino¹ · Franz Nogoy¹ · 김준기¹ · 소윤섭¹ · 조용구^{1*}

¹충북대학교 농업생명환경대학 식물자원학과, ²국립종자원

Classification of Korean Rice Varieties Based on Agro-morphological Traits

Me-Sun Kim¹, Hye-Jung Lee^{1,2}, Dal-A Yu¹, Jae-Young Song¹, Marjohn Nino¹,
Franz Nogoy¹, Joonki Kim¹, Yoon-Sup So¹, Yong-Gu Cho^{1*}

¹Department of Crop Science, Chungbuk National University, Cheongju 28644, Korea

²Current Address, Korea Seed & Variety Service, Kimchon 39660, Korea

Abstract : The International Union for the Protection of New Varieties of Plants (UPOV) promotes an effective system of plant variety protection and encourages the development of new varieties of plants. This international convention was initiated to standardize the system efforts and strengthen policy. The establishment of cultivar discrimination system is very important to distinguish varieties between domestic and foreign agricultural products. It is necessary for the protection of breeders' rights. In addition, it will help for more efficient and quality management of plant breeding. This study was conducted to identify and group rice varieties based on agro-morphological characteristics such as plant height, panicle length, number of tillers, culm length, leaf length, leaf width, leaf pigments and flag leaf angles. Using these parameters, statistical analysis classified a total of 243 rice varieties bred in Korea into four groups. Most rice varieties did not exhibit anthocyanin pigments on the leaves particularly on the first leaf, leaf blade, leaf sheath and auricle, except for varieties classified as black rice. Results of phylogenetic and principal component analysis (PCA) indicated that these varieties formed three largely distinct clusters according to their ecotype and morphological differentiation. This result would be useful in rice varietal identification for the protection of breeders' variety rights.

Keywords : Korean rice variety, Morphological character, Cluster analysis, Principal component analysis

서 언

우리나라에서 벼 품종의 육종은 1950년대에 교배육종을 통하여 다수성 품종 육성에 주력하면서 내도복성과 내병성 증진에 노력해왔고, 1970년대에는 japonica와 indica 벼의 원연교잡으로 다수성과 내도복성 및 내병충성 증대에 모든 육종적 노력을 집중하는 한편, 통일형 품종의 결점인 내냉성, 내탈립성, 그리고 미질개선을 위해 노력해왔다. 1980년대의 큰 냉해 피해 이후에는 내냉성 강화에 힘을 기울였고, 쌀의 과잉생산과 소비자들의 양질미 선호도가 증가함에 따라 쌀의 미질개선에 힘 기울였다(Lee et al. 2012). 그 후에는 japonica 벼 품종의 초형 개선에

의한 수량성과 안정성 증대에 주력하였다. 그리고, 쌀의 소비증 국내 육성 벼 243 품종의 형태적 특성에 따른 품종 분류대 및 부가가치 증진을 위한 가공적성 증진, 병충해 및 기상재해 복합저항성의 유전적 보완이 강화되었다. 1990년대 이후에는 쌀의 품질이 우수하고 밥맛이 좋은 양질미의 수량성과 안정성 증대를 위해 노력함과 동시에 가공용 벼 품종개발 등 품종의 다양화에 주력하여 왔다(Kang et al. 2004, Breseghello 2013). 최근에는 WTO와 FTA에 의하여 쌀 시장의 개방에 대비하여 국제경쟁력 강화를 위한 고품질 품종육성에 주력하면서 남북통일에 대비하고 식량안보의 확보를 위한 쌀 수량성 증대연구와 병해충과 재해에 복합저항성을 지닌 품종개발을 지속하고 있다(Kim 2013).

벼 육종가가 새로운 품종을 육성하기 위해서는 10~15년의 기간이 소요되는 등 많은 시간과 노력이 투여되는 반면, 일단

*Corresponding author (E-mail: ygcho@cbnu.ac.kr, Tel: +82-43-261-2514, Fax: +82-43-273-2242)

(Received on July 8, 2016. Accepted on August 8, 2016.)

새로운 품종이 육성되어 공개되면 많은 경우 다른 사람들에게 의해 쉽게 재생산되어 그것을 개발한 육종가에게 투자에 대한 적절한 보상의 기회를 박탈해 버림으로써 의욕을 상실하게 만든다(Choi et al. 2004). 따라서 식물육종 분야에 있어서 개인 육종가, 발명자에게 그 신품종의 개발에 이르기까지의 시간과 노력, 재정적인 면 등에 있어서 그 대가로서 지식재산권을 부여하여 보호할 필요가 있는 것이다(Park et al. 2009). 이 밖에도 종자에 대해 지식재산권을 부여함으로써 정부 또는 공공부분의 역할을 줄이고 민간부분을 통해 품종육성 및 종자생산을 할 수 있다는 점 등을 지적하기도 한다.

1998년부터 도입된 품종보호제도는 식물 신품종 육성자의 권리를 법적으로 보장하여 주는 지적재산권의 한 형태로 육종자에게 배타적인 상업적 독점권을 부여하여 품종 개발을 촉진하기 위한 제도이다(Seong et al. 2009).

품종보호제도의 규범화는 신품종의 육성을 촉진하고 우수품종의 이용을 확대하여 농업의 생산성을 증대시킬 수 있는 긍정적인 효과가 기대되며, 이와 같이 법적으로 식물품종에 대한 보호를 의무화하려는 추세가 전세계적으로 확대되고 있다(Choi 2004, Garris et al. 2005).

최근 선진국들은 신품종 종자산업을 국가경쟁력의 새로운 원천으로 인식하여 지원을 강화하고 있다(Osteen et al. 2012, SCBD 2005). 그러나 우리나라는 ‘종자산업법’에서 신품종 육성자의 권리를 법적으로 보장하기 위해 식물 신품종보호제도를 채택하고 있으나 품종보호에 관한 국내에서의 논의는 활발하지 않다. 또한, 식물에 대한 국내 유통품종 중 법정종자를 제외한 품종의 특성 데이터베이스를 구축하여 관리하는 노력이 부족하며, 품종 간 유전적 최소거리(Minimum Distance)에 대한 법정 분쟁 및 환경과 외형의 차이에 따른 특성별 차이에 대한 연구가 미흡하다. 벼에서는 1984년부터 1994년까지 수많은 형태적 돌연변이 특성들의 연관군지도를 작성하여 그들의 염색체별 위치

를 보고한 바 있는데(Kinoshita 1984~1994, Causse et. al. 1994), 이러한 형태적 특성의 변이를 이용하면 수많은 유사한 벼의 품종을 구분하는 데에 매우 효과적으로 이용될 수 있을 것으로 생각된다.

우리나라 국가기관에서는 18 개 작물에 대한 DNA Profile의 데이터베이스를 구축하여 다방면에 활용하고 있으나, 국제신품종보호동맹 (International Union for the Protection of New Varieties of Plants, UPOV)의 사례와 같이 국가적이고 표준화된 데이터베이스의 구축은 미흡한 실정으로 차후에 분석결과의 신뢰도 및 공정성에 대한 문제점이 발생할 수 있을 것으로 예상된다.

따라서, 본 연구에서는 우리나라에서 품종으로 육성되어 유통되고 있는 벼 품종의 외형의 차이에 따른 품종별 형태적 특성을 농촌진흥청 및 국립종자원의 조사기준에 따라서 조사하였고(RDA 2003a) 벼 품종 간 유전적 거리를 탐색하여 각 품종의 형질 측정을 통해 데이터베이스를 구축하여 표준품종을 선정하고 품종의 형태적 특성뿐만 아니라 DNA profiling에 의하여 품종에 대한 구분체계를 확립할 수 있는 기초 정보를 제공하고자 하였다.

본 보고에서는 국내에서 육성하여 국립종자원을 통하여 보급되고 있는 243개 벼 품종에 대하여 특히 조사가 가능한 벼의 형태적 특성을 질적형질과 양적형질 모두를 세밀하게 조사하여 주성분 분석 및 군집 분석을 수행하였고 그를 통하여 품종군을 분류하였으며, 초형, 이삭 및 현미의 모양을 사진을 찍어 직접 비교에 활용할 수 있게 제공하였다.

재료 및 방법

시험재료 및 재배법

시험재료는 국내에서 육성된 벼 품종 중 1982년부터 2013년까지 국립종자원을 통하여 보급된 243개 벼 품종이 사용되었다

Table 1. List of 243 rice varieties bred in Korea from 1982 to 2013.

No.	Variety	Type (J/T) ^z	Year	Cross Combination
1	Keumyeong	J	2009	Sanbaek/Iksan423//Sangju22
2	Durujinmi	J	2010	Suwon478/Odae
3	Manho	J	2002	Singeumo/Iksan423
4	Sandeuljinmi	J	2006	Sambaek/Yongseong12
5	Samcheonbyeon	J	1995	Unbong/Fukei126
6	Unjangbyeon	J	1994	Unbong//Suwon244/Hwasetoramoggi//Sangnam
7	Wonpyeong	J	1999	Hwaseong Mutant
8	Jeokjinju	J	2000	Obong//Obong/Syare

9	Josaengheukchal	J	2004	Dongbukna149/Sx864
10	Joan	J	2003	Jinmi/jinbu10
11	Jungmo1011	J	2010	Suwon478/Unbong33
12	Jinbong	J	2000	Jinmi/Unbong
13	Taebong	J	2000	SR13330-13-3-5-2-1/Jinbu10
14	Taeseong	J	2002	Cheolwon49/Jinbu10
15	Hwanggeumbora	J	2006	Jinbu/Odae//Fukei126
16	Heukseonchal	J	1999	Sanghaehyanghyeolra Mutant
17	Geurubyeyo	J	1997	Suwon313/Cheolwon42
18	Geumseong	J	2003	Hajjeubosi/Hyangmi
19	Dunnaebyeo	J	1992	SR11139-48-1/Suwon320
20	Manna	J	2005	Iksan438/Ilmi
21	Manchu	J	2001	Jinmi/Unbong12
22	Sambaekbyeoy	J	1993	Gosihikkari/TR2406-2-1-1//Hukkurikku115/Cheolwon29
23	Sangsanbyeoy	J	1993	Sobaek/Daeseong
24	Sangjuchalbyeoy	J	1997	YR4117-99-1-1-2-4/ YR4200-2-3-2-2
25	Seolbaek	J	2010	Suwon360/Geuru
26	Sinunbong1	J	2005	Sangju/Unbong17
27	Ansanbyeoy	J	1995	Rax102-123/Seonam
28	Wonhae	J	2007	Dongjin Mutant
29	Wolbaek	J	2010	Milkikwin/Geuru
30	Inwol	J	1998	Fukei127/Unbong
31	Joun	J	2009	SR14880-173-3-3-2-2-2/Unbong20
32	Jungmo1007	J	2009	Sanbaek/Unbong19
33	Junghwabyeyo	J	1995	Etznal126/Bokgang//Daeseong
34	Cheongbaekchal	J	2010	SR20261-5-4-10/Jinbupal
35	Hyangmibyeoy	T	1996	IR841-76-1/Suwon334
36	Goun	J	2004	Jinbu10/Jinbu17
37	Namil	J	2002	Ilpum/Namyang7
38	Manan	J	1998	Milyang110/Milyang110/Yeongdeok7
39	Munjang	J	1999	Sangsan/Suwon397
40	Sangmi	J	1998	Sambaek/Ou316
41	Saasangju	J	2001	Junghwa/Sambaek
42	Odae1	J	2005	Ilpum/Jinbu10
43	Undu	J	1998	Odae/Jinbu13
44	Wonpum	J	2004	Ilpum Mutant
45	Jeokjinjuchal	J	2010	Sangnambat/Jeokjinju
46	Jopyeong	J	2010	HR16683-46-3/HR18129-B-16
47	Daechan	J	2009	Nongan/2*Suwon403
48	Jungmo1010	J	2009	SR18390-9-7-2-5/IR66160-5-2-3-2
49	Heukjinjubyeyo	J	1997	Yongjeong4/Segeum
50	Jakwangchal	J	2007	Jinbupal/Jagwangdo
51	Hwanggeumchal	J	2007	Sakita/Jinbupal
52	Daeanyeyo	J	1994	Oseto/Seomjin
53	Seomyeong	J	2009	Kyehwa21/Milyang165
54	Seojin	J	1996	Aiji37/Sangpung
55	Juanbyeoy	J	1994	Seolak/Gosihikkari//Samnam
56	Hojin	J	2000	Hwayeong//Dongjin/Milyang95
57	Hwarang	J	2003	Iksan420/YR13616Acpl
58	Hwamyeyongbyeoy	J	1997	Hwayeong/SR14779-HB234-32
59	Nampyeongbyeoy	J	1997	HR11735-82-1-5-5/Milyang95
60	Naepungbyeoy	J	1995	Milyang97//OU316/Iri373

61	Dasan	T	1995	Suwon332/Suwon333
62	Dasan1	T	2006	Bangal//Yongju/Dasan
63	Daesanbyeo	J	1996	Milyang95/Suwon366
64	Dongbo	J	2010	Yeongdeok19/Gosihikkari
65	Dongan	J	1996	Milyang95/HR5119-12-1-5
66	Dongjin2	J	2005	Milyang165/Iksan438//HR14018-B-1-1/Iksan435
67	Dongjinchal	J	1998	SR11155-4-2/Doyonisiki//Milyang95
68	Donghaejinmi	J	2006	Milyang64/4*Milyang165
69	Malgeumi	J	2006	Hwayeong/Aichi 76
70	Mikwang	J	2009	SR15926-10-2-3-3-3/Iksan431
71	Mihyang	J	1999	Seomjin/Dongbuk144//IR392
72	Baekjinju	J	2001	Ilpum Mutant
73	Boramchan	J	2009	HR21124-B-59/Kyehwa24
74	Saekyehwa	J	2001	Ilpum//Mangeum/Jyukeoi830
75	Seonong10	J	2005	Hwacheong Mutant
76	Seonong16	J	2010	Seonong8/Saengdongchal
77	Segyejinmi	T	2009	Milyang160/Yongju
78	Sujin	J	1999	Milyang95//Milyang96/Milyang106
79	Woncheong	J	2003	Chucheong Mutant
80	Junam	J	2000	Hwayeong//Sangju/Ilpum
81	Jungmo1013	J	2010	Iksan469//Dongjin1/Iksan474
82	Cheonga	J	2006	Ginuhikari/Yangju
83	Cheonghaejinmi	J	2009	Samjiyeon/SR14694-57-4-2-1-3-2-2//Iri402
84	Chinnong	J	2010	Iksan450/YR21258-GH3
85	Keunseom	T	2006	Milyang165*3/Gosihikari
86	Pyeongang	J	2003	Iksan438/HR15003-69-B-3
87	Hyangmibyeo1	T	1993	IR841-76-1/Suwon334
88	Hoan	J	1998	Kanto149/Milyang95
89	Hwasin1	J	2005	HR13185-B-B-100//Iri407/HR14732-B
90	Heuknam	J	1997	Tamjin/Sanghaehyanghyeolna
91	Gancheokbyeo	J	1992	Aichi37/Seomjin
92	Gangbaek	J	2006	Suwon345/DV85
93	Gangchan	J	2010	Iksan421/Suwon438
94	Goami2	J	2002	Ilpum Mutant
95	Goami4	J	2009	Suwon464/Daeon
96	Gopumbyeo	J	2004	SR15225-B-22-1-2//54BC-68/Iri369
97	Geumnambyeo	J	1994	Jinju/Suwon346
98	Geumbyeol	J	1995	Chukri830//KantoPL5//Milyang79/Aichi65
99	Namcheonbyeo	T	1995	YR3299-34-2-2/Suwon318
100	Nokwonchalbyeo	J	2007	Nokmi Mutant
101	Nongan	J	1993	SR5204-30-2/Pungsan
102	Dami	J	2006	Iksan438/Iksan426
103	Dasan2	T	2009	Suwon450/SR21356
104	Daejin	J	1996	Daeseong/Seomjin
105	Deurimi2	J	2009	SNDH-39/Junam
106	Manpung	J	2000	Nakdong//Iri390/Milyang111
107	Boseokchal	J	2004	Hwayeong//Tamjin/2*Sinseonchal
108	Samkwang	J	2003	Suwon361/Milyang101
109	Samdeok	J	2002	YR/12733-B-B-5-1/Milyang101
110	Sanggal	J	2007	Ilpum Mutant
111	Sangok	J	2003	Milyang101/YR8697 Acp 19
112	Seonong12	J	2006	Nampung//EM40/Nampung*2
113	Seonong6	J	2002	Hwacheong Mutant

114	Seonong8	J	2005	Hwacheong Mutant
115	Seonong9	J	2006	Hwacheong Mutant
116	Seoan1	J	2005	Namyang9/Kyehwa7
117	Seoeun	J	2008	Hwacheong Mutant
118	Seopyeong	J	2003	Hwayeong/HR11752
119	Seokjeong	J	2001	Namyang7/SR11340-30-4-1-3-2
120	Seolgaeng	J	2001	Ilpum Mutant
121	Sura	J	1998	Suwon345/KantoPL 4//Suwon345
122	Sinmyeongheukchal	J	2006	Heuknam/Milyang153
123	Sinbaek	J	2010	Iksan469/HR23966-22-1-2
124	Sintoheukmi	J	2008	Heuknam/Milyang153
125	Areum	T	1999	YR3299-34-2-2/Suwon318
126	Anda	T	1998	SR11532-4/SR14502F2
127	Anmi	J	2010	Junam/IR65482-7-216-1-2//3*Junam
128	Yangjobyeo	J	1994	HR7874-AC77/HR8140/AC59
129	Yeonghaebyeo	J	1997	Milyang101/Chucheong
130	Orae	J	2007	Ilpum Mutant
No.	Variety	Type (J/T)	Year	Cross Combination
131	Onnuri	J	2005	Milyang165/HR14732-B-67-2-3
132	Wonkwang	J	2000	Seomjin Mutant
133	Wonmi	J	2000	Seomjin Mutant
134	Wonhwang	J	1998	Milyang96/Iri390/Milyang5
135	Yunnongchamssal	J	2008	Dong poor rice/Glutinous rice
136	Joryeongbyeo	J	1992	Gosihikari/Seonam//Milyang79
137	Jongnam	J	2001	Milyang96/YR12734-B-B-22-2
138	Juan1	J	2005	Ilpum/SR18392-HB683-104
139	Jungmo1001	J	2004	Cheolwon52/SR21049
140	Jungmo1012	J	2007	Sambaek/Ou329
141	Jungmo1014	J	2010	Milyang207/Samgwang
142	Jungmo1009	J	2009	YR22207//Iksan2443/Milyang165
143	Jungsan	J	2000	Sambaek/Milyang107
144	Cheongdam	J	2006	SR19200-HB826-34/Juan
145	Cheongho	J	2004	Iri407/Iri417
146	Pungmi	J	2004	Milyang165//YR16365Acp13/Iksan438
147	Pungmi1	J	2005	YR13616Acp1/Milyang122
148	Hangangchal1	T	2006	Hangangchal/YR8208-20
149	Hanmaeum	J	2004	Milyang165//YR16365Acp13/Iksan438
150	Hanareum2	T	2010	Milyang181/Milyang154
151	Haepyeong	J	2000	Milyang101/Akichikara
152	Haepyeongchal	J	2004	Haepeong Natural Mutant
153	Hyangnambyeo	J	1995	Iri389/Dohoku144
154	Hongjinju	J	2006	SR18164F2/Suwon383
155	Hwanambyeo	J	1993	Milyang95/Tamjin
156	Hwadongbyeo	J	1997	Daejin/SR13345-20-1
157	Hwabong	J	1998	Milyang95/Iri390//Milyang101/Iri390
158	Hwasambyeo	J	1996	Milyang101/Iri389
159	Hwaseonchalbyeo	J	1992	Inabawase/Dongjin
160	Hwasinbyeo	J	1995	Iri390/Milyang110
161	Hwaan	J	2000	Suwon362/SR10778-2-2
162	Hwajungbyeo	J	1993	SaSanisiki/Cheonma
163	Heukgawichal	J	2008	Iksan427/Gawichal
164	Geonganghongmi	J	2010	Sugary*2/Milyang152

165	Goami	J	2000	Milyang95//Kimcheon poor rice/Ilpum
166	Geuman	J	2002	SR11878-14-4-1/Suwon345
167	Geumobyeo2	J	1996	Milyang96//YR6419Acp13/Palgong
168	Namgangbyeo	J	1997	.Milyang95/Milyang96
169	Nogyang	T	2006	Yongmun/IR67396-16-3-3-1
170	Daeripbyeol	J	1993	Niheunmasari ms/BG29
171	Daepyeong	J	2002	HR14028-AC-5/Milyang122
172	Dongwon	J	2011	-
173	Manmi	J	2002	Milyang95//Bukryukbanna/Milyang95
174	Mogyang	J	2010	SR24592-HB2319/IR3165-B-6-1-1
175	Milkikwin	J	2007	Gosihikari Mutant
176	Sampyeong	J	2000	Suwon345/SR11340-46-5-4-1
177	Saechucheong BIL-3	J	1999	Daeseong/5*Chucheong//Bonggwang/5*Chucheong
178	Seonong15	J	2010	Heukjinju/Floury rice
179	Seolhyangchal	J	1999	Miyagaori/2*Suwon357
180	Seongjochal	J	2008	Wangchal Mutant
181	Suan	J	2010	HR20017-B-19-3-1/Iksan467
182	Sindongjin	J	1999	Hwayeong/YR13604ACP22
183	Aranghyangchalbyeol	J	1997	Sinseonchal/Tohoku144
184	Onojjinoijji	J		-
185	Wonchu	J	2001	Chucheong Mutant
186	Jungan	J	1999	Namyang7/2*Hapcheon1
187	Jinbo	J	2009	Yeongdeok26/Gosihikari
188	Jinsang	J	2008	Yumetsukushi/Milkykwin
189	Garak	J	2009	Chucheong/Sinseonchal
190	Kwangan	J	1998	Namyang7/SR14779-HB234-31
191	Geumtap	J	2008	Sinseonchal//Juan/쌀사레
192	Noreunjachal	J	2008	Sinseonchal//Juan/쌀사레
193	Nunbora	J	2006	Iksan433/Miyadanamojji
194	Daeripjami	J	2009	C3GHi/Daerip1
195	Dongjin1	J	2001	Hwayeong/HR12800-AC21
196	Manwol	J	2001	Milyang120/Hwayeong
197	Manjong	J	2009	Yeongdeok34/Nampeong
198	Seonong14	J	2008	Heukjinju/Geodae
199	Sobi	J	1999	Hwayeong/YR13604ACP22
200	Yeongan	J	2001	Milyang122/YR13616 Acp1
201	Jinpum	J	1999	SR14703-60-5-GH1/Suwon353
202	Keunnun	J	2005	Ilpum Mutant
203	Keunnunjami	J	2007	Heukjinju/Suwon425
204	Hopyeong	J	2003	Hitobebore/Hwajin
205	Heukkwang	J	2003	Hwajin//Gilrimheukmi/Ilpum
206	Nongho	J	1998	Chukei1016/Milyang79
207	Mipum	J	2010	Gosihikari//KyeHwa21/Junam
208	Syupojami	J	2009	C3GHi/EM76
209	Suryeojinmi	J	2010	Milyang182/SR21144-17-3-2
210	Heukhyang	J	2000	Mangeum/SX864
211	Baekjinju1	J	2005	Ilpum(MNU)-10-2-GH1-3/Seoan
212	Jungmo1015	J	2010	Sindongjin/Milkikwin
213	Saenuri	J	2007	Kyehwa/HR14026-B-68- 6-1-5
214	Yeonghojinmi	J	2009	Hiddeumeaborea/Junam
215	Hiami	J	2009	Jinmi Mutant
216	Dacheong	J	2008	Iksan450/YR21258-GH3

217	Daeyabyeo	J	1992	Gwandong PL5/Gwandong PL3//Daechong
218	Deuraechan	J	2008	Milyang165//Sindongjin/YR19105Acp222
219	Boseok	J	2008	Ginuhikari//HR19621AC6/Sobi
220	Seogan	J	2002	Hr11752-11-14-4/HR10213 -11-3-5
221	Unmi	J	2007	Samcheon/HR17870
222	Jungmo1008	J	2010	Iksan451/Sangju21
223	Jinbaek	J	2008	HR15204-38-3//Milyang165/Iksan438
224	Honong	J	2009	Unbong31 Mutant
225	Hwanggeumnodeul	J	2007	Milyang165/HR15151-B-21-3
226	Honggwang	J	2009	Chucheong/Sinseonchal
227	Goami3	J	2008	Suwon464/Daeon
228	Danmi	J	2009	Sugary/Seomjin
229	Mogwoo	J	2009	Dasan//Suwon431/IR71190-45-2-1
230	Baekokchal	J	2009	Dongjinchal/YR19700
231	Boseokheukchal	J	2008	SR18638-B=B-B-18-2/H31
232	Joami	J	2008	Yukara/Tonggae112//Sambaek
233	Jungmo1004	J	2007	Sugary/HwayeongAcp33
234	Jungmo1005	J	2009	Milyang165//Seomjin//Iksan438/Sangju13
235	Jungmo1006	J	2009	Samgwang/Nakdong
236	Jinsumi	J	2008	Milyang165//YR16195-B-B-B-21-1/Milyang169
237	Cheongcheongjinmi	T	2009	IR401/Ilpum
238	Cheongpungheukchal	J	2009	Sindongjin/Heukseonchal
239	Pyeongwon	J	2007	Jinbu19/Sanjiyeon4
240	Handeul	J	2007	Sangju/Tomoemasari//Geuru
241	Haeoreumi	J	2008	Milyang165/Haepyeong
242	Hoban	J	2007	Hidomebore/Jinbu
243	Cheongnam	J	2009	Ginuhikari/Milyang189

^ZJ: *Japonica* type, T: *Tongil* type

(Table 1). 벼 품종은 2013년 국내 육성 벼 243 품종의 형태적 특성에 따른 품종 분류와 2014년에 충북대학교 시험포장에 이양하여 관찰, 조사하였다. 품종 당 1주 1본씩, 16주/월, 3열로 이양하였으며 물 관리와 비료의 사용 및 병, 해충의 방제는 농촌진흥청 벼 표준재배법에 준하여 수행하였다(RDA 2003a).

조사항목

벼의 형태적 특성 조사를 위하여 양 끝에 심겨진 3개체씩을 배제하고 안쪽에 심겨진 10주를 기준으로 하여 3반복으로 농촌진흥청 조사기준 및 국립종자원에서 규정한 벼 표준조사방법에 따라 조사하였다(RDA 2003a). 형태적 특성 조사를 크게 잎, 줄기, 출수기로 나누어 조사하였으며, 질적형질은 육안조사를 실시하였고, 양적형질은 1000 mm 금속 자를 이용하여 측량 조사를 실시하였다(Table 2). 조사항목은 초장, 간장, 제3엽기에 제1엽의 안토시아닌 색소 유무, 잎의 녹색정도 및 안토시아닌 색소 유무, 잎혀의 모양 및 색, 잎몸의 길이 및 너비를 조사하였으며, 잎몸의 길이 및 너비는 출수 직후에 상위 3번째 엽의

길이를 측정하였다. 줄기는 지엽의 자세 및 굵기, 길이, 모양을 조사하였으며, 이때 지엽의 자세는 지엽의 만곡성 정도를 가지고 조사하였고 줄기의 굵기는 주간 엽신의 폭이 가장 넓은 부위를 조사하였다. 출수기는 조사개체의 50%가 출수하였을 때를 출수기로 하였다.

통계분석

통계프로그램 SAS version 9.2 (SAS Institute Inc.)과 Multibase program (<http://www.numericaldynamics.com>)을 이용하였으며 2013년도와 2014년도에 243개 품종으로부터 얻어진 데이터를 분산분석을 통해 유의성을 검정하였으며, 벼 243개 품종들간의 형태적 특성에 기초한 품종군을 분류하고자 시각적으로 가장 두드러지는 형태적 특성인 출수기를 비롯하여 초장, 간장, 수장, 분얼수 및 잎몸의 너비와 길이 등 총 7개의 형질을 이용하여 주성분 분석과 군집분석을 수행하였다.

Table 2. Agronomic traits investigated in the morphological analysis of rice varieties.

Characteristics	Subsection	Stage
Qualitative trait		
1 st leaf	Color	Nursery stage
Leaf	Intensity of green color Distribution of anthocyanin	Maximum tiller number stage
Ligule	Color, Shape	Maximum tiller number stage
Flag leaf	Attitude	Heading stage, Yellow ripe stage
Culm	Thickness, Attitude	Dough ripe stage, Maturity stage
Quantitative trait		
Leaf blade	Length, Width (cm)	Maximum tiller number stage
Culm	Length (cm)	Dough ripe stage, Maturity stage
Time of heading	Time of heading (Day)	50% heading of rice
Panicle	Panicles per plant (No.) Length (cm)	Dough ripe stage, Maturity stage

결과 및 고찰

형태적 특성

(1) 잎의 길이와 너비

벼 243 품종의 잎 형태적 특성을 비교 분석하기 위하여 잎몸의 길이 및 너비를 조사하였다.

잎몸의 길이 및 너비는 출수 직후에 상위 3번째 엽의 길이를 측정하였으며 개화시기별로 나타낸 결과는 Table 3과 같다. 잎몸 길이에서 조생종은 월백이 23.44 cm 국내 육성 벼 243 품종의 형태적 특성에 따른 품종 분류로 가장 짧았고 중모1012가 47.88 cm로 가장 길었으며 평균은 30.96 cm를 나타내었다. 중생종은 큰섬이 21.76 cm로 가장 짧았고 소비가 59.10 cm로 가장 길었으

며 평균은 39.74 cm로 나타났으며, 만생종은 흑향이 40.6 cm로 가장 짧으며 백진주1호가 56.44 cm로 가장 길었고 평균은 46.98 cm를 나타내었다.

잎몸 너비는 조생종에서 운장벼가 0.86 cm로 가장 좁게 나타났고 흑가위찰이 1.74 cm로 가장 넓은 잎몸 너비를 가진 것으로 나타났고 평균은 1.27 cm로 관찰되었다. 중생종은 금남벼가 1.02 cm로 최소치를, 성조찰이 2.54 cm로 최대치를 나타냈고 평균은 1.38 cm로 나타났으며, 만생종은 신토흑미가 1.08 cm로 가장 좁게 나타났고 목우가 1.80 cm로 넓게 나타났으며 평균은 1.46 cm로 세 품종군 중에서 가장 넓게 관찰되었다.

출수기에 따른 잎의 길이는 출수기에 따라 유의한 차이가 있는 것으로 나타나서 만생종, 중생종, 조생종 순으로 컸으며, 잎의 너비 또한 출수기에 따라 유의한 차이가 있는 것으로 인정되

Table 3. Variation of leaf lengths and widths in three groups of rice varieties in terms of flowering time.

Characteristic	Min	Max	Mean	SD	F	p-value
Leaf length(cm)					53.67***	<0.001
Early ^Z	23.44	47.88	30.96	±4.55		
Medium	21.76	59.10	39.74	±7.18		
Late	40.6	56.44	46.98	±6.08		
Leaf width(cm)					9.12***	<0.001
Early	0.86	1.74	1.27	±0.17		
Medium	1.02	2.54	1.38	±0.20		
Late	1.08	1.8	1.46	±0.26		

^ZEarly = Early flowering rice varieties; Medium = Medium flowering rice varieties; Late = Late flowering rice varieties. * P < 0.05, ** P < 0.01, *** P < 0.001, ns: not significant.

Table 4. The investigation of existence of rice varieties showing anthocyanin pigmentation in 1st leaf, leaf blade, and ligule.

Distribution of anthocyanin	No. of variety		
	1 st leaf	Leaf bade	Ligule
Absent	235	239	237
Present	8	4	6

* Variety name were provided in Supplementary Table 1, 2, and 3.

Table 5. Number of rice varieties with different green color intensity of leaf blades and with ligule

Green color density	No. of variety	Shape (Ligule)	No. of variety
Pale green	2	Partedness	235
Medium green	182	Sharp	8
Dark green	59		

* Variety name were provided in Supplementary Table 4 and 5.

어 만생종, 중생종, 조생종 순으로 컸다(Table 3). 만생종에서 잎몸의 길이와 너비의 평균 최고치가 관찰되었으며 이 같은 결과는 엽면적 및 엽신건물중의 증가로 식물체 전체의 광이용율 및 광합성 효율을 극대화시켜 동화물질의 축적에 크게 기여할 것으로 생각된다.

(2) 잎의 색과 형태

벼 243 품종의 잎의 형태적 특성을 비교 분석하기 위하여 유묘기의 제1엽색, 잎의 녹색 정도 및 안토시아닌 유무, 잎혀의 모양과 색을 조사하였다.

파종 후 3~4엽기에 품종별로 제1엽의 부위별 안토시아닌색소의 유무를 조사한 결과는 Table 4와 같다. 제1엽색이 녹색을 띠는 품종은 동진벼를 포함하여 235개 품종에서 관찰되었고, 자색을 띠는 품종은 자광찰, 신명흑찰, 흑찰, 품미1호, 눈보라, 서은, 조생흑찰, 보석흑찰, 신토흑미 총 9개 품종으로 조사되었다. 또한, 자색을 나타내는 부위가 신명흑찰, 신토흑미, 조생흑찰과 같이 서로 다른 위치에서 부분적으로 나타남을 관찰할 수 있었다(Table 4, Supplementary Table 1).

잎과 잎혀에서 안토시아닌색소의 유무를 조사한 결과 주로 흑미로 분류되는 조생흑찰, 보석흑찰, 신토흑미 등에서 나타났고 대부분의 벼 품종에서는 나타나지 않았다(Table 4, Supplementary Table 2, 3).

파종 후 3~4엽기에 품종별로 제1엽색 및 잎의 형태를 조사한 결과는 Table 5와 같다. 벼 품종의 최고분얼기에 잎의 녹색정도와 잎에 포함된 안토시아닌색소의 유무를 조사하였고, 잎의 녹색정도는 연녹색, 중간녹색, 진녹색으로 분류하여 관찰하였다. 조

사결과 잎이 연녹색을 나타내는 품종은 고품, 품미1호로 전체 품종의 0.82%를 차지하였고, 중간녹색은 182개 품종으로 75.31%, 진녹색은 59개 품종으로 23.87%의 비율로 나타났다(Table 5, Supplementary Table 4).

뽕죽한 형태의 잎혀를 가진 품종은 만추, 문장, 화랑, 평안, 미품, 흑선찰, 성조찰, 홍진주 총 8개 품종으로 나타났고, 235개 품종은 갈라진 잎혀의 모양으로 나타났다(Table 5, Supplementary Table 4).

(3) 줄기의 길이 및 분얼수

벼 243개 품종의 줄기의 형태적 특성은 길이 및 분얼수로 나누어 특성 비교를 수행하였다.

이삭의 길이를 제외한 간장과 분얼수를 개화시기별로 분석한 결과는 Table 6과 같다. 조생종은 설백이 62.42 cm로 가장 작았고 자광찰이 105.42 cm로 가장 컸으며 평균 76.86 cm로 나타났고, 중생종은 주안1호가 60.71 cm로 가장 작았고 서농8호가 84.60 cm로 가장 컸으며 평균 78.20 cm로 3가지 품종군 중에서 가장 크게 나타났다. 만생종은 흑향이 61.16 cm로 가장 작았고 목우가 84.60 cm로 가장 컸으나 평균은 74.32 cm로 가장 작게 나타났다.

벼의 분얼수는 조생종에서 대찬과 황금보라가 6.8개로 가장 적었고 보석이 16.0개로 가장 많았으며 평균 10.9개로 나타났다. 중생종은 목양이 5.6개로 243개 품종 중 최소치로 나타났고 보석찰이 18.8개로 243개 품종 중 최대치를 나타냈으며 평균 11.0개로 품종군 중 가장 높게 나타났다. 만생종은 목우가 7.4개로 가장 적었고 서농16이 10.4개로 가장 많았으며 평균은 8.9개

Table 6. Variation of culm lengths and panicles per plants in three rice varieties groups in terms of flowering time.

Characteristics	Min	Max	Mean	SD	F	p-value
Culm length(cm)					1.62 ^{ns}	0.20
Early ^Z	62.42	105.42	76.86	±6.62		
Medium	60.72	104.60	78.20	±8.07		
Late	61.16	84.60	74.32	±7.93		
Panicles per plant(no.)					3.84 [*]	0.02
Early	6.8	16.0	10.9	±1.97		
Medium	5.6	18.8	11.0	±2.28		
Late	7.4	10.4	8.9	±1.15		

^ZEarly = Early flowering rice varieties; Medium = Medium flowering rice varieties; Late = Late flowering rice varieties,

* P < 0.05, ** P < 0.01, *** P < 0.001, ns: not significant.

Table 7. Number of flag leaf angle at heading stage and yellow ripe stage in 243 rice varieties.

Flag leaf attitude	No. of variety	
	Heading stage	Yellow ripe stage
Erect	222	190
Semierect	21	45
Medium	0	8

* Variety name were provided in Supplementary Table 6 and 7.

로 나타났다(Table 6).

출수기에 따른 간장의 길이는 출수기 간에 유의성이 나타나지 않았으나, 분얼수의 분석 결과 출수기에 따라 유의한 차이가 있는 것으로 인정되어 중생종이 11개로 가장 많았고 조생종(10.9 개), 만생종(8.9개) 순이었다(Table 6).

(4) 줄기의 형태와 색

벼 243개 품종의 줄기의 정성적 특성은 출수기 및 황숙기 지엽의 자세, 줄기 굵기 및 자세로 나누어 특성 비교를 수행하였다.

지엽의 경우 출수기와 황숙기로 나누어 조사를 하였으며, 이삭을 기준으로 지엽의 각도가 0-20°는 직립형, 35-50°는 반직립형, 50°이상의 각도를 보이는 것은 수평형으로 나누어 구분하였다.

그 결과 출수기에는 222개 품종이 직립형으로 나타났고, 품미 1호, 두루진미, 운장벼, 오대1호벼 등 36개 품종이 반직립형으로 나타났으며, 수평형을 나타내는 품종은 없었다(Table 7), 황숙기에는 190개 품종이 직립형, 45개 품종이 반직립형으로 나타났고, 오대1호, 큰눈자미, 청백찰, 홍광, 운두, 대립벼1호, 흑진주, 호반 총 8개 품종이 수평형을 나타내었다(Table 7). 일부 벼 품종을 제외하고는 출수기와 황숙기에 동일한 형태를 띠었으며 대체로

직립형이 많았고, 수평은 극히 일부에 불과하였다. 가장 대표적인 수평형 품종은 큰눈자미이며, 반직립형 품종은 삼천벼로 나타났다.

벼의 호숙기 및 성숙기에 줄기 모양을 비교한 결과, 205개 품종이 직립형을 나타냈고, 설백, 남일, 만호, 조아미를 포함한 총 8개 품종이 보통 개형, 29개 품종이 반직립형으로 대부분의 벼 품종이 직립의 형태를 보였다(Table 9). 줄기의 굵기는 24개 품종이 가는 굵기, 164개 품종이 보통 굵기, 55품종이 굵은 줄기로 대부분이 보통의 줄기 굵기를 나타내었다(Table 8).

(5) 초장 및 수장

벼 243개 품종을 대상으로 초장 및 수장을 조사하였으며 개화 시기별로 분석한 결과는 Table 9와 같다.

초장의 경우 조생종은 조운이 87.00 cm로 가장 작았고 자광찰이 135.80 cm로 가장 컸으며 평균 99.80 cm로 나타났고, 중생종은 호진벼가 91.20 cm로 가장 작았고 서농14호가 138.10 cm로 가장 컸으며 평균 107.70 cm로 3가지 품종군 중에서 가장 크게 나타났다. 만생종은 흑향이 85.60 cm로 가장 작았고 목우가 128.10 cm로 가장 컸으나 평균은 103.40 cm로 나타났다.

벼의 이삭길이는 조생종에서 대진벼가 17.42 cm로 가장 짧았

Table 8. Number of culm thickness and attitude in 243 rice varieties.

Thickness	No. of variety	Attitude	No. of variety
Thin	24	Erect	205
Medium	164	Semierect	29
Thick	55	Medium	8

* Variety name were provided in Supplementary Table 8 and 9.

Table 9. Variation of plant height and panicle length in three rice variety groups in terms of flowering time.

Characteristics	Min	Max	Mean	SD	F	p-value
Plant height(cm)					19.55***	<0.001
Early	87.0	135.8	99.8	±7.95		
Medium	91.2	138.1	107.7	±8.96		
Late	85.6	128.1	103.4	±13.38		
Panicle length(cm)					15.18***	<0.001
Early	17.42	25.80	21.51	±1.68		
Medium	15.74	29.64	20.02	±2.19		
Late	16.66	21.70	18.92	±1.82		

^zEarly = Early flowering rice varieties; Medium = Medium flowering rice varieties; Late = Late flowering rice varieties,

* P < 0.05, ** P < 0.01, *** P < 0.001, ns: not significant.

Table 10. Eigenvalues and cumulative proportion of the first 7 principal components in 243 rice varieties.

Principal component	Eigen value	Difference	Proportion	Cumulative
1	2.22194685	0.48697278	0.3174	0.3174
2	1.73497407	0.499865537	0.2479	0.5653
3	1.2351087	0.45826783	0.1764	0.7417
4	0.77684087	0.2008556	0.111	0.8527
5	0.57598527	0.23700687	0.0823	0.935
6	0.3389784	0.22281256	0.0484	0.9834
7	0.11616584		0.0166	1

고 흑진주벼가 25.80 cm로 가장 길었으며 평균 21.57 cm로 품종군 중 가장 높게 나타났다. 중생종은 삼광벼가 15.74 cm로 최소치로 나타났고 목양이 29.64 cm로 243개 품종 중 최대치를 나타냈으며 평균 20.02 cm로 나타났다. 만생종은 슈퍼자미가 16.66 cm로 가장 짧았고 목우가 21.70 cm로 가장 길었으며 평균은 18.92 cm로 나타났다(Table 9).

출수기에 따른 초장의 길이는 출수기에 따라 유의한 차이가 있는 것으로 나타나서 중생종, 만생종, 조생종 순으로 컸으며, 이삭의 길이 또한 출수기에 따라 유의한 차이가 있는 것으로 인정되어 조생종, 중생종, 만생종 순으로 컸다(Table 7).

출수기 및 형태형질 관련 특성

출수기는 조사 개체의 50%가 출수하였을 때를 기준으로 하여 조사하였으며, 이앙 후 일수에 따라 구분하였다. 이앙 후 70일 이전에 출수한 품종은 조생종, 70일에서 90일 사이에 출수한 품종은 중생종, 90일 이후에 출수한 품종은 만생종으로 분류하였다(Fig. 1A). 또한, 고품질 벼 품종 재배면적 확대와 품질 고급화 실현을 위해 국내 육성 벼 243 품종을 대상으로 벼 품종별 키, 수형, 이삭 및 종자의 모양과 크기 등 특성을 표시해 농업인들이 올바른 품종 선택으로 고품질 쌀을 안전하게 생산할 수 있도록 정보 제공 및 품종에 대한 구분체계를 확립하여 육종의 기초 자료로 활용하고자 하였다 (Fig. 1B, Supplementary Figure 1).

Fig. 1. A, Phenotype of early, medium and late flowering rice varieties at 70 days after transplanting. (A) Boseok, (B) Sinunbong1, (C) Daechan (D) Cheongbaekchal,, (E) Suryeojinmi. B, Morphological traits, plant type, panicle shape and grain size in three representative varieties for early-, medium-, and late-ecotype, respectively, among 243 Korean rice varieties (See Supplementary Fig. 1 for more varieties).

다변량 분석에 의한 품종 분류

(1) 주성분 분석

공시된 243개 벼 품종의 개별적 특성이 객관성이 있게 평가되어야 하나 단일 변량에 의하여 전체적인 변이의 구조를 파악하기는 어렵다. 따라서 다변량 분석법을 조사형질보다 적은 수의 주성분으로 요약한 후 여러 변수들의 구조적 관계를 파악하는데 사용하였다(Choi 2011).

벼 243개 품종을 대상으로 출수기, 초장, 간장, 수장, 분얼수, 잎몸너비 및 잎몸길이 총 7가지 형질을 이용하여 통계프로그램 SAS 9.2의 PRINCOMP procedure로 주성분분석을 수행하였

다. 주성분분석으로 고유치와 각 주성분의 기여도를 나타낸 결과는 Table 10과 같다.

분석결과 고유치와 기여도에서 각각 제1주성분은 고유치가 2.2로 나타나며 전체 분산 중 31.7%를 설명할 수 있으며, 제2주성분은 1.73에 24.8%, 제3주성분은 1.24에 17.6%, 제4주성분은 0.78에 11.1%, 제5주성분은 0.58에 8.2%, 제6주성분은 0.34에 4.8%, 제7성분은 0.12에 1.7%로 각각 설명할 수 있었다. 이를 전부 해석하지 않고 이들 중 자료의 변동을 많이 설명하는 몇 개의 주성분만을 선택하는데 일반적으로 고유 값이 1이상, 또는 누적 분산비율이 80% 이상이 되도록 주성분의 수를 결정한다(RDA 2003b). 본 실험은 고유 값 1이상을 기준으로 3개의 주성분을 사용하여 전체자료 변동의 74%를 설명 할 수 있었는데,

이는 3개의 주성분을 통해 7개 형질에서 측정된 전체 분산의 74%를 설명할 수 있어 벼 243개 품종의 출수기 및 주요 수량구성 요소에 따른 분류가 가능하다는 것을 알 수 있었다. 또한, 이들 주성분이 함축하고 있는 형태적 형질 중 어떤 형질이 이들 3가지 주성분에 요약되어 있는지를 유추하고 시각적으로 분류하여 집단의 경향을 알아보기 위하여 Multibase program을 이용하여 주성분분석을 수행하였다. 농업적으로 중요한 7가지 형태적 형질 값을 이용하여 품종들의 구조적 관계를 확인한 결과는 Fig. 2와 같다. 제1주성분은 전체변이의 27.1%, 제2주성분은 10%, 제3주성분은 7%를 설명할 수 있었다. 주성분분석 결과 얻어진 243개 품종의 분류적 관계는 품종간의 뚜렷한 집구현상은 관찰되지 않았지만, 조생종(Red group), 중생종(Yellow group) 및 만생종(Blue group) 크게 3개의 집단으로 구분되었다(Fig. 2B). 이를 바탕으로, 외부형태학적 형질을 조사한 결과 파종 후 개화까지의 소요일수, 잎의 길이 및 너비에 따라 국내 육성 벼 품종의 분류에 의미가 있는 것으로 확인되었다(Fig. 2A). 또한, 국내 육성 벼 품종을 3개의 집단으로 구분하는 형태 특성 중 출수기가 큰 비중을 차지하는 것을 알 수 있다. 이러한 결과는 타작물의 형태적 특성 즉, 식물크기, 과실의 길이 및 너비 등 차이에 의한 분류 및 유연관계를 분석한 Huh et al.(2008)의 결과처럼 작물의 형태적 형질이 품종간의 분류가 의미가 있음을 보여주었다. 그러나 정확한 품종군 분류를 위해서는 보다 많은 형태적 형질의 이용과 분자학적 접근이 필요할 것으로 생각된다.

(2) 군집 분석

군집분석은 주어진 관찰치 중에서 유사한 것들을 몇몇 집단으로 그룹화하여 각 집단의 성격을 파악함으로써 데이터 전체의 구조에 대한 이해를 돕고자 하는 분석방법으로 주성분분석 자체로도 정보를 얻을 수 있지만 주성분분석은 차후 분석을 위한 준비단계로 변수의 수를 줄여 그래프를 통한 관찰치의 위치를 시각화하여 집단의 경향을 파악하는데 사용한다(Choi 2011).

벼 243개 품종의 7개 형질을 이용하여 주성분분석을 수행한 후 그래프상의 품종들의 배열로 분류관계를 파악하고자 하였으며, 전체 변이의 74%를 설명할 수 있는 제3주성분까지의 주성분 점수를 이용하여 군집분석을 실시한 결과는 Fig. 3과 같다.

군집분석을 수행한 결과에 따라서 품종을 분류는 Table 11와 같으며, 0.9에서 4개 군으로 분류되었다. 제I군은 벼 82개 품종이 속하였는데, 그 중 조생종 품종이 60개, 중생종 품종이 22개로 나타났다. 전체적으로 잎몸의 길이 및 너비가 다른 군에 비해 길고 짧지만 많은 분얼수를 가지는 것으로 나타났고, 특이적으로

조생종인 홍광의 출수기가 이앙 후 40일로 월등히 빠른 것으로 나타났다. 제II군은 중생종 품종 114개, 만생종 품종 2개 등 총 116개의 품종이 속하였는데, 다른 군에 비해 잎몸의 길이 및 너비가 긴 특성을 보였다.

제I군과 제II군에 속하는 품종이 243개 품종 중 81%의 비율을 차지하며 제I 군에는 조생종 품종이 큰 비율을 차지하고 있으며 제II 군에는 중생종 품종이 크게 나타났으며 이것으로 품종을 구분하는 표현형 중 큰 비중을 차지하는 형태적 특성이 출수기임을 알 수 있다. 각 군에 속해 있는 품종을 육종 및 인공교배에 이용하고자 한다면 출수가 늦은 품종은 파종을 앞당기고 출수가 빠른 품종은 파종을 늦추어 양친의 개화기를 일치시킴으로써 교잡을 유도할 수 있을 것으로 여겨진다.

제III군은 중생종 24개 품종, 만생종 7개 품종이 속하였으며, III군의 31개 품종은 다른 품종들에 비해 초장과 간장의 길이가 긴 특성을 보이고 있었다. 제IV군은 조생종 품종 6개, 중생종 품종 8개 총 14개 품종이 속하였다. 분얼수는 많지 않으나 초장과 간장의 길이가 월등히 길었으며, 중생종인 성조찰의 잎몸 너비가 2.54cm로 가장 넓게 나타났다. 제III군과 제IV군은 엽면적 및 엽신건물중의 증가로 식물체 전체의 광이용율 및 광합성 효율을 극대화시켜 동화물질의 축적에 크게 기여할 것으로 생각된다.

본 연구에서는 국내 243개 벼 품종에 대하여 형태적 변이의 범위를 이해하고, 이들의 효율적 관리와 유용한 신품종의 발굴을 위한 기초연구로서 형태적 특성조사 결과를 바탕으로 주성분 분석 및 군집 분석을 수행하였다. 출수기를 포함한 초장, 간장, 수장, 분얼수, 잎몸너비, 잎몸길이 등 7가지 특성을 이용하여 군집분석을 실시한 결과, 총 4개의 품종군으로 분류되었다.

제I 군과 제II 군은 출수기가 중요한 표현형으로 나타났으며 제III군과 제IV군은 특이적으로 우월한 표현형 위주의 분류군으로 나타났다. 주성분분석과 군집분석 결과 크게 4개 군집, 조생종, 중생종 및 만생종, 그리고 혼합된 군집 경향으로 분류되어 동일한 결과를 보였다. 품종간의 뚜렷한 군집 현상이 관찰되지 않고, 한 개의 군집내에 숙기에 따른 많은 품종들이 속해 있는 결과는 서로 간에 유사한 표현형을 가지고 있는 유전적 유사도가 높은 교배친이 교배모본으로 이용된 데서 기인한 것이라 생각된다(Lee et al. 2015).

따라서 본 연구 결과로 고찰해 볼 때 신품종 발굴을 위해서는 육종의 초기세대에서 숙기형태별로 계통을 구분하여 적응성을 평가하는 것이 바람직할 것으로 판단된다.

Fig. 2. Principal component analysis (PCA) on the seven morphological traits of the 243 Korean rice varieties. PC 1 and PC 2 refer to the first and second principal components, respectively.

적 요

국내 243개 벼 품종에 대하여 잎과 줄기의 형태적 특성조사 결과를 바탕으로 주성분 분석 및 군집 분석을 수행하였다. 잎몸의

길이 및 너비는 만생종에서 각각 46.98 cm과 1.46 cm로 가장 길고 넓게 나타났으며, 흑미로 분류되는 품종의 제1엽, 본엽, 잎혀에서 안토시아닌 색소를 나타내었다. 잎의 녹색정도는 중간 녹색을 가진 품종의 비율이 높았고 갈라진 잎혀의 모양을 가진

Fig 3. Cluster dendrogram generated by flowering time and phenotypic characters for the 243 rice accessions. *Red: early flowering variety, Yellow: medium flowering variety, Blue: late flowering variety.

품종이 대부분이었다. 또한, 간장의 길이는 조생종에서 최대치 105.42 cm가 나타났고 중생종에서 최소치 60.72 cm가 나타났으며, 분얼수는 중생종에서 최소치 5.6개와 최대치 18.8개가 모두 나타났다. 일부 벼 품종을 제외하고는 출수기와 황숙기 지엽 자세가 동일한 형태를 띠었으며 대체로 직립형이 많았고, 수평은 극히 일부에 불과하였다. 출수기를 포함한 정량적인 7가지 특성을 이용하여 군집분석을 실시한 결과, 총 4개의 품종군으로 분류되어 주성분분석과 동일한 결과, 조생종, 중생종 및 만생종, 그리고 혼합된 군집 경향을 보였다. 이와 같은 결과, 신품종 발굴을 위해서는 육종의 초기세대에서 숙기형태별로 계통을 구분하여 적응성을 평가하는 것이 바람직할 것으로 판단된다.

사 사

본 연구결과는 농림축산식품부의 재원으로 농림수산식품기술기획평가원의 농생명산업기술개발사업의 지원(313043-03-3-HD030)을 받아 연구되었으며 이에 대한 감사를 표합니다.

Supplementary Information

아래와 같은 보충자료는 한국육종학회지 홈페이지 (www.koreabreedjournal.org)에서 확인할 수 있습니다.

Supplementary Table 1. The investigation of existence of rice variety showing anthocyanin pigmentation in 1st leaf.

Supplementary Table 2. The investigation of existence of rice variety showing anthocyanin pigmentation in leaf blade.

Supplementary Table 3. The investigation of existence of rice variety showing anthocyanin pigmentation in ligule.

Supplementary Table 4. Frequency distribution of rice variety showing different green color intensity in leaf blade.

Supplementary Table 5. Frequency distribution of rice variety showing types in ligule.

Supplementary Table 6. Frequency distribution of flag leaf angle in 243 rice varieties at heading stage.

Supplementary Table 7. Frequency distribution of flag leaf angle in 243 rice varieties at yellow ripe stage.

Supplementary Table 8. Frequency distribution of culm thickness in 243 rice varieties.

Table 11. Varietal composition of each group classified by the cluster analysis using the first four principal component scores.

Group	No.	Maturity	Variety name
I	82	Early ^Z (60)	Geumyeong, Durujinmi, Boseok, Dunnae, Sandeuljinmi, Inwol, Wolbaek, Sambaek, Heukseonchal, Naepungbyeol, Joryeongbyeol, Geurubyeol, Sangsanbyeol, Manna, Sangjuchalbyeol, Sinunbong1, Munjang, Daejin, Seolbaek, Joun, Manho, Josaengheukchal, Saesangju, Jungmo1001, Manan, Jungmo1012, Samcheonbyeol, Handeul, Unjangbyeol, Junghwabyeol, Wonpyeong, Jinbong, Geumseong, Unmi, Joan, Hwadongbyeol, Jopyeong, Jungmo1008, Sangmi, Joami, Jeokjinju, Jungmo1010, Wonpum, Taeseong, Honong, Jungmo1011, Manchu, Goun, Taebong, Ansanbyeol, Wonhae, Jungmo1007, Hyangmibyeol2, Pyeongwon, Cheongbaekchal, Odae1, Undu, Hoban, Heukgawichal, Honggwang
		Medium (22)	Namcheonbyeol, Hyangmibyeol1, Segyejinmi, Anda, Keunseom, Hanareum2, Geumobyeol1, Dasan2ho, Cheongdam, Gangbaek, Daeanbyeol, Jungsan, Cheonga, Seonong12, Pungmi, Seolhyangchal, Manwol, Dasan, Juanbyeol, Areum, Orae, Dasan1
II	116	Medium (114)	Seomyeong, Donghaejinmi, Gopum, Yangjobyeol, Sanggol, Chinnong, Daeyabyeol, Mipum, Geumnambyeol, Deurimi2, Samgwang, Dongjin2, Cheongpungheukchal, Saenuri, Goami2, Goami4, Juan1, Jinbaek, Anmi, Jungmo1006, Jinsumi, Cheongnam, Seojin, Hoan, Daesanbyeol, Dongan, Mihyang, Yeonghaebyeol, Wongwang, Sangok, Heukgwang, Dongwon, Hwamyongbyeol, Sura, Dongbo, Manpung, Hwabong, Jungan, Wongcheong, Gangchan, Seonong15, Jinsang, Cheongho, Haepyeong, Hojin, Boramchan, Jongnam, Haiami, Suan, Jinbo, Saekyehwa, Jungmo1013, Hanmaeum, Suryeojinmi, Samdeok, Sinbaek, Hwarang, Sinmyeongheukchal, Onnuri, Wonhwang, Deuraechan, Nampyeongbyeol, Junam, Onojjinoijji, Seopyeong, Hwanggeumnodeul, Sampyeong, Jungmo1005, Seolgaeng, Yeongan, Haeoreumi, Sindongjin, Dongjinchal, Baekjinju, Seogan, Nongho, Malgeumi, Cheonghaejinmi, Pyeongan, Dami, Seonong10, Aranghyangchalbyeol, Sujin, Hwasambyeol, Anmi, Hongjinju, Cheongcheongjinmi, Haepyeongchal, Hwaan, Daeripbyeol1, Jinpum, Jungmo1009, Hyangnambyeol, Pungmi1, Hwanambyeol, Daepyeong, Goami3, Manjong, Manmi, Gwangan, Boseokheukchal, Jungmo1004, Migawng, Wonchu, Seoan1, Gancheokbyeol, Heuknam, Seonong9, Seoeun, Seonong6, Boseokchal, Nokwonchal, Seonong8, Saechucheong BIL-3
		Late (2)	Jungmo1015, Syupeojami
		Medium (24)	Namgangbyeol, Keunnunjami, Wonmi, Milkikwin, Hwaseonchalbyeol, Nokyang, Hwasinbyeol, Noreunjachal, Hwasin1, Jungmo1013, Goami, Sobi, Seokjeong, Hwajungbyeol, Geumobyeol2, Hopyeong, Daeripjami, Baekokchal, Dongjin1, Geonganghongmi, Garak, Geuman, Seonong14, Geumtap
III	31	Late (7)	Seonong16, Sintoheukmi, Heukhyang, Yeonghojinmi, Baekjinju1, Dacheong, Mogwoo
		Early (6)	Hwanggeumbora, Namil, Jeokjinjuchal, Heukjinjubyel, Hwanggeumchal, Daechan
		Medium (8)	Hangangchal1ho, Keunnunjami, Nongan, Yunnongchamssal, Mokyang, Nunbora, Jagwangchal, Seongjochal
IV	14		

^ZEarly = Early flowering rice varieties; Medium = Medium flowering rice varieties; Late = Late flowering rice varieties

Supplementary Table 9. Frequency distribution of culm attitude at maturity in 243 rice varieties.

Supplementary Figure 1. Morphological traits, plant type, shape and size of panicle and grain in 243 Korean rice varieties.

REFERENCES

- Bresegheello F. 2013. Traditional and Modern Plant Breeding Methods with Examples in Rice (*Oryza sativa* L.). J. Agric. Food Chem 61(35):8277-8286.
- Causse MA, Fulton TM, Cho YG, Ahn SN, Chunwongse J, Wu K, Xiao J, Yu Z, Ronald PC, Harrington SE. 1994. Saturated molecular map of the rice genome based on an interspecific backcross population. Genetics 138(4):1251-74.
- Choi KJ. 2004. Recent Issues Discussing about Plant Variety Protection. Korean J. Breed. Sci 36(1):76-93.
- Choi HG. 2011. Assessment of Major Physicochemical Components on Waxy Corn Hybrids *Zea mays* L.). Ph.D. Diss., Chungnam Nat'l Univ., Daejeon, Korea.

5. Choi KJ, Kim CH. 2010. Impact of Introduction of Plant Breeders' Right. The Journal of Intellectual Property 5(3):124-126.
6. Garris AJ, Tai TH, Coburn J, Kresovich S, McCouch S. 2005. Genetic Structure and Diversity in *Oryza sativa* L. Genetics 169:1631-1638.
7. Huh YC, Solmaz I, Sari N. 2008. Morphological characterization of Korean and Turkish watermelon germplasm. Proceedings of the IXth EUCARPIA Meeting on Genetics and Breeding of Cucurbitaceae, INRA, Avignon (France), pp 327-334.
8. Kang KH, Seo JP, Cho YC, Choi IS, Ahn SN, Hwang HG, Choi HC, Choi SH, Moon HP. 2004. Use of wild rices for rice improvement in Korea. Korean J. Breed. Sci 36(1):104-109.
9. Kim DI. 2013. Improving National Implementing Law of WTO Government Procurement Agreement and FTA Government Procurement Chapter. Korea Legislation Research Institute.
10. Kinoshita T. 1984-1994. Report of the committee on gene symbolization and linkage map. Rice Genet. Newsl. 1-11.
11. Lee GY, So YS, Kim HS, Hong JH, Jang EK, Lee KH, Yun GS, Hwang SG. 2015. Variation of Growth and Yield Component Characters and Varietal Classification by Multivariate Analysis in Korean Soybean Varieties. Korean J. Int. Agric 27(3):348-357.
12. Lee AS, Cho YS, Kim IJ, Ham JK, Jang JS. 2012. The Quality and Yield of Early Maturing Rice Varieties affected by Cultural Practices in Gangwon Plain Region. Korean J. Crop Sci 57(3):233-237.
13. Osteen CD, Gottlieb J, Vasavada U. 2012. Agricultural Resources and Environmental Indicators, 2012 Edition. U.S. Department of Agriculture. EIB-98.
14. Park JH. 2009. Research on the Protection Plans of a seed-related invention. Korean Intellectual Property Office.
15. Rural Development Administration (RDA). 2003a. Standard methods for agricultural experimental experiments.
16. Rural Development Administration (RDA). 2003b. Agricultural science technology research analysis base.
17. Secretariat of the Convention on Biological Diversity. 2005. Handbook of the Convention on Biological Diversity Including its Cartagena Protocol on Biosafety, 3rd edition.
18. Seong MH, Park HT, Han HS, Park SC. 2009. Evaluation and Strategies for Protecting the plant variety. Korea Rural Economic Institute 6(7)

Supplementary Table 1. The investigation of existence of rice variety showing anthocyanin pigmentation in 1st leaf.

Anthocyanin (1 st leaf)	No. of variety
Absent	Orae, Hwanggeumchal, Heukgawichal, Geumseong, Joryeongbyeo, Hwanambyeo, Sambaekbyeo, Sangsanbyeo, Junghwabyeo, Namcheonbyeo, Naepungbyeo, Geumobyeo1, Hwasambyeo, Daesanbyeo, Geumobyeo2, Namgangbyeo, Aranghyangchalbyeo, Yeonghaebyeo, Sangjuchalbyeo, Gancheokbyeo, Geumnambyeo, Yangjobyeo, Unjangbyeo, Hwasinbyeo, Hyangnambyeo, Samcheonbyeo, Dongan, Nampyeongbyeo, Hwaseonchalbyeo, Dunnaebyeo, Hwajungbyeo, Daeripbyeo1, Nongan, Hyangmibyeo1, Juanbyeo, Daeanbyeo, Dasan, Ansanbyeo, Daejin, Seojin, Hyangmibyeo2, Hwadongbyeo, Geurubyeo, Hwamyongbyeo, Heukjinjubyo, Sura, Gwangan, Anda, Undu, Mihyang, Dongjinchal, Hoan, Inwol, Hwabong, Nongho, Manan, Wonhwang, Sangmi, Jungan, Seolhyangchal, Sobi, Sindongjin, Areum, Munjang, Sujin, Sampyeong, Taebong, Hwaan, Jinbong, Jeokjinju, Hojin, Manpung, Heukhyang, Junam, Haepyeong, Jungsan, Goami, Jinpum, Saekyehwa, Manchu, Saesangju, Jongnam, Manwol, Yongan, Seolgaeng, Baekjinju, Seokjeong, Samdeok, Manmi, Geuman, Namil, Taeseong, Daepyeong, Manho, Goami2, Hwarang, Sangok, Saechucheong BIL-3, Dongjin1, Samgwang, Joan, Heukgwang, Hopyeong, Seopyeong, Pyeongan, Cheongho, Boseokchal, Hanmaeum, Pungmi, Haepyeongchal, Goun, Gopum, Hwasin1, Sinunbong1, Onnuri, Manna, Dongjin2, Odae1, Juan1, Keunnun, Seon1, Baekjinju1, Gangbaek, Nokyang, Dami, Dasan1, Donghaejinmi, Malgeumi, Sandeuljinmi, Cheogndam, Cheonga, Keunseom, Hangangchal1, Hongjinju, Hwanggeumbora, Milkikwin, Seongjochal, Noreunjachal, Geumtap, Seonong6, Seonong10, Seonong8, Seonong9, Seonong12, Sanggol, Wonpyeong, Wongwang, Wonmi, Heukseonchal, Woncheong, Wonpum, Nokwonchal, Wonhae, Jungmo1012, Seonong16, Jinsang, Anmi, Segyejinmi, Manjong, Durujinmi, Daeripjami, Jeokjinjuchal, Suryeojinmi, Dongwon, Jungmo1007, Garak, Chinnong, Jopyeong, Geonganghongmi, Sinbaek Onojjinoijji, Dongbo, Mipum, Wolbaek, Seolbaek, Migwang, Cheonghaejinmi, Jungmo1001, Geumyeong, Boramchan, Seomyeong, Yoonnongchamssal, Hanareum2, Gangchan, Deurimi2, Syupeojami, Keunnunjami, Jungmo1010, Daechan, Dasan2, Mokyang, Joun, Seonong15, Jinbo, Goami4, Cheongbaekchal, Suan, Seonong14, Jungmo1009, Jungmo1014, Jungmo1011, Jungmo1015, Jungmo1013, Saenuri, Yeonghojinmi, Hiam, Dacheong, Daeyabyeo, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Honggwang, Goami3, Danmi, Mogwoo, Baekokchal, Joami, Jungmo1004, Jungmo1005, Jungmo1006, Jinsumi, Cheongcheongjinmi, Cheongpungheukchal, Pyeongwon, Handeul, Haeoreumi, Hoban, Cheongnam
Present	Jagwangchal, Sinmyeongheukchal, Heuknam, Pungmi1, Nunbora, Seoeeun, Josaengheukchal, Boseokheukchal, Sintoheukmi

Supplementary Table 2. The investigation of existence of rice variety showing anthocyanin pigmentation in leaf blade.

Anthocyanin (Leaf blade)	No. of variety
Absent	<p>Orae, Hwanggeumchal, Heukgawichal, Geumseong, Joryeongbyeo, Hwanambyeo, Sambaekbyeo, Sangsanbyeo, Junghwabyeo, Namcheonbyeo, Naepungbyeo, Geumobyeo1, Hwasambyeo, Daesanbyeo, Geumobyeo2, Namgangbyeo, Aranghyangchalbyeo, Yonghaebyeo, Sangjuchalbyeo, Gancheokbyeo, Geumnambyeo, Yangjobyeo, Unjangbyeo, Hwasinbyeo, Hyangnambyeo, Samcheonbyeo, Dongan, Nampyeongbyeo, Hwaseonchalbyeo, Dunnaebyeo, Hwajungbyeo, Daeripbyeo1, Nongan, Hyangmibyeo1, Juanbyeo, Daeanyeo, Dasan, Ansanbyeo, Daejin, Seojin, Hyangmibyeo2, Hwadongbyeo, Geurubyeo, Hwamyongbyeo, Heukjinjubyo, Sura, Gwangan, Anda, Undu, Mihyang, Dongjinchal, Hoan, Inwol, Hwabong, Nongho, Manan, Wonhwang, Sangmi, Jungan, Seolhyangchal, Sobi, Sindongjin, Areum, Munjang, Sujin, Sampyeong, Taebong, Hwaan, Jinbong, Jeokjinju, Hojin, Manpung, Heukhyang, Junam, Haepyeong, Jungsan, Goami, Jinpum, Saekyehwa, Manchu, Saesangju, Jongnam, Manwol, Yongan, Seolgaeng, Baekjinju, Seokjeong, Samdeok, Manmi, Geuman, Namil, Taeseong, Daepyeong, Manho, Goami2, Hwarang, Sangok, Saechucheong BIL-3, Dongjin1, Samgwang, Joan, Heukgwang, Hopyeong, Seopyeong, Pyeongan, Cheongho, Boseokchal, Hanmaeum, Pungmi, Haepyeongchal, Goun, Gopum, Hwasin1, Sinunbong1, Onnuri, Manna, Dongjin2, Odae1, Juan1, Keunnun, Seoan1, Baekjinju1, Gangbaek, Nokyang, Dami, Dasan1, Donghaejinmi, Malgeumi, Sandeuljinmi, Cheogndam, Cheonga, Keunseom, Hangangchal1, Hongjinju, Hwanggeumbora, Milkikwin, Seongjochal, Noreunjachal, Geumtap, Seonong6, Seonong10, Seonong8, Seonong9, Seonong12, Sanggol, Wonpyeong, Wongwang, Wonmi, Heukseonchal, Woncheong, Wonpum, Nokwonchal, Wonhae, Jungmo1012, Seonong16, Jinsang, Anmi, Segyejinmi, Manjong, Durujinmi, Daeripjami, Jeokjinjuchal, Suryeojinmi, Dongwon, Jungmo1007, Garak, Chinnong, Jopyeong, Geonganghongmi, Sinbaek Onojjinoljiji, Dongbo, Mipum, Wolbaek, Seolbaek, Migwang, Cheonghaejinmi, Jungmo1001, Geumyeong, Boramchan, Seomyeong, Yoonnongchamssal, Hanareum2, Gangchan, Deurimi2, Syupeojami, Keunnunjami, Jungmo1010, Daechan, Dasan2, Mokyang, Joun, Seonong15, Jinbo, Goami4, Cheongbaekchal, Suan, Seonong14, Jungmo1009, Jungmo1014, Jungmo1011, Jungmo1015, Jungmo1013, Saenuri, Yeonghojinmi, Hiami, Dacheong, Daeyabyeo, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Honggwang, Goami3, Danmi, Mogwoo, Baekokchal, Joami, Jungmo1004, Jungmo1005, Jungmo1006, Jinsumi, Cheongcheongjinmi, Cheongpungheukchal, Pyeongwon, Handeul, Haeoreumi, Hoban, Cheongnam, Jagwangchal, Sinmyeongheukchal, Heuknam, Pungmi1, Nunbora</p>
Present	<p>Seoeun, Josaengheukchal, Boseokheukchal, Sintoheukmi</p>

Supplementary Table 3. The investigation of existence of rice variety showing anthocyanin pigmentation in ligule.

Anthocyanin (Ligule)	No. of variety
Absent	Geumyeong, Durujinmi, Manho, Unjangbyeo, Wonpyeong, Joan, Jungmo1011, Taebong, Taeseong, Geurubyeo, Manna, Manchu, Sangsanbyeo, Seolbaek, Sinunbong1, Ansanbyeo, Wolbaek, Inwol, Joun, Jungmo1007, Junghwabyeo, Goun, Namil, Manan, Munjang, Saesangju, Odae1, Undu, Wonpum, Jopyeong, Daechan, Jungmo1010, Daesanbyeo, Seomyeong, Seojin, Juanbye, Hojin, Hwarang, Nampyeongbye, Daesanbye, Dongbo, Dongan, Dongjin2, Dongjinchal, Donghaejinmi, Malgeumi, Migwang, Mihyang, Baekjinju, Boramchan, Saekyehwa, Seonong10, Sujin, Woncheong, Junam, Jungmo1013, Cheonga, Cheonghaejinmi, Pyeongan, Hoan, Hwasin1, Gancheokbye, Gangbaek, Gangchan, Goami2, Goami4, Geumnabye, Geumobye, Nongan, Dami, Daejin, Deurimi2, Manpung, Boseokchal, Samgwang, Samdeok, Sanggol, Sanggol, Sangok, Seonong6, Seonong8, Seonong9, Seonong1, Seooun, Seopyeong, Seokjeong, Seolgaeng, Sura, Sinbaek, Anmi, Yangjobye, Yeonghaebye, Onnuri, Wongwang, Wonmi, Wonhwang, Jongnam, Juan1, Jungmo1012, Jungmo1014, Jungmo1009, Jungsan, Cheongho, Hanmaeum, Hanmaeum2, Haepyeong, Haepyeongchal, Hyangnambye, Hwabong, Hwasambye, Hwasinbye, Hwaan, Hwajungbye, Geonganghongmi, Goami, Geuman, Geumobye2, Namgangbye, Nokyang, Daeripbye1, Daepyeong, Dongwon, Manmi, Mokyang, Milkikwin, Sampyeong, Saechucheong BIL-3, Seonong15, Seolhyangchal, Suan, Sindongjin, Aranghyangchalbye, Onojjinoijji, Wonchu, Jungan, Jinbo, Jinsang, Garak, Gwangan, Noreunjachal, Daeripjami, Dongjin1, Manjong, Seonong14, Sobi, Yeongan, Jinpum, Keunnun, Keunnunjami, Hopyeong, Heukgwang, Nongho, Mipum, Syupeojami, Juryeojinmi, Heukhyang, Baekjinju1, Jungmo1015, Saenuri, Hiami, Dacheong, Daeyabye, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Honggwang, Goami3, Baekokchal, Joami, Jungmo1004, Jungmo1006, Jinsumi, Cheongcheongjinmi, Pyeongwon, Handeul, Haeoreumi, Hoban, Cheongnam, Gopum, Pungmi1, Sandeuljinmi, Samcheonbye, Jeokjinju, Jinbong, Hwanggeumbora, Heukseonchal, Geumseong, Dunnaebye, Sambaekbye, Sangjuchalbye, Wonhae, Cheongbaekchal, Hyangmibye2, Sangmi, Jeokjinjuchal, Heukjinjuby, Jagwangchal, Hwanggeumchal, Hwamyongbye, Naepungbye, Dasan, Dasan1, Seonong16, Segyejinmi, Chinnong, Keunseom, Hyangmibye1, Namcheonbye, Nokwonchal, Dasan2, Seonong12, Areum, Anda, Orae, Yoonnongchamssal, Joryeongbye, Jungmo1001, Cheongdam, Pungmi, Hangangchal1, Hongjinju, Hwanambye, Hwadongbye, Hwaseonchalbye, Seongjochal, Geumtap, Nunbora, Manwol, Yeonghojinmi, Danmi, Mogwoo, Jungmo1005, Cheongpungheukchal
Present	Heukgawichal, Sinmyeongheukchal, Josaengheukchal, Boseokheukchal, Heuknam, Sintoheukmi

Supplementary Table 4. Frequency distribution of rice variety showing different green color intensity in leaf blade.

Green color density	No. of variety
Pale green	Gopum, Pungmil
Medium green	Geumyeong, Durujinmi, Manho, Unjangbyeon, Wonpyeong, Joan, Jungmo1011, Taebong, Taeseong, Geurubyeo, Manna, Manchun, Sangsanbyeon, Seolbaek, Sinunbong1, Ansanbyeon, Wolbaek, Inwol, Joun, Jungmo1007, Junghwabyeon, Goun, Namil, Manan, Munjang, Saesangju, Odae1, Undu, Wompum, Jopyeong, Daechan, Jungmo1010, Daeanbyeon, Seomyeong, Seojin, Juanbyeon, Hojin, Hwarang, Nampyeongbyeon, Daesanbyeon, Dongbo, Dongan, Dongjin2, Dongjinchal, Donghaejinmi, Malgeumi, Migwang, Mihyang, Baekjinju, Boramchan, Saekyehwa, Seonong10, Sujin, Woncheong, Junam, Jungmo1013, Cheonga, Cheonghaejinmi, Pyeongan, Hoan, Hwasin1, Gancheokbyeon, Gangbaek, Gangchan, Goami2, Goami4, Geumnambyeon, Geumbyeon, Nongan, Dami, Daejin, Deurimi2, Manpung, Boseokchal, Samgwang, Samdeok, Sanggol, Sanggol, Sangok, Seonong6, Seonong8, Seonong9, Seonong1, Seoeun, Seopyeong, Seokjeong, Seolgaeng, Sura, Sinbaek, Anmi, Yangjobyeon, Yeonghaebyeon, Onnuri, Wongwang, Wonmi, Wonhwang, Jongnam, Juan1, Jungmo1012, Jungmo1014, Jungmo1009, Jungsan, Cheongho, Hanmaeum, Hanmaeum2, Haepyeong, Haepyeongchal, Hyangnambyeon, Hwabong, Hwasambyeon, Hwasinbyeon, Hwaan, Hwajungbyeon, Geonganghongmi, Goami, Geuman, Geumbyeon2, Namgangbyeon, Nokyang, Daeripbyeon1, Daepyeong, Dongwon, Manmi, Mokyang, Milkikwin, Sampyeong, Saechucheong BIL-3, Seonong15, Seolhyangchal, Suan, Sindongjin, Aranghyangchalbyeon, Onojjinoijji, Wonchu, Jungan, Jinbo, Jinsang, Garak, Gwangan, Noreunjachal, Daeripjami, Dongjin1, Manjong, Seonong14, Sobi, Yeongan, Jinpum, Keunnun, Keunnunjami, Hopyeong, Heukgwang, Nongho, Mipum, Syupeojami, Juryeojinmi, Heukhyang, Baekjinju1, Jungmo1015, Saenuri, Hiami, Dacheong, Daeyabyeon, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Honggwang, Goami3, Baekokchal, Joami, Jungmo1004, Jungmo1006, Jinsumi, Cheongcheongjinmi, Pyeongwon, Handeul, Haoreumi, Hoban, Cheongnam
Dark green	Sandeuljinmi, Samcheonbyeon, Jeokjinju, Josaengheukchal, Jinbong, Hwanggeumbora, Heukseonchal, Geumseong, Dunnaebyeon, Sambaekbyeon, Sangjuchalbyeon, Wonhae, Cheongbaekchal, Hyangmibyeon2, Sangmi, Jeokjinjuchal, Heukjinjubyon, Jagwangchal, Hwanggeumchal, Hwamyongbyeon, Naepungbyeon, Dasan, Dasan1, Seonong16, Segyejinmi, Chinnong, Keunseom, Hyangmibyeon1, Heuknam, Namcheonbyeon, Nokwonchal, Dasan2, Seonong12, Sinmyeongheukchal, Sintoheukmi, Areum, Anda, Orae, Yoonnongchamssal, Joryeongbyeon, Jungmo1001, Cheongdam, Pungmi, Hangangchal1, Hongjinju, Hwanambyeon, Hwadongbyeon, Hwaseonchalbyeon, Heukgawichal, Seongjochal, Geumtap, Nunbora, Manwol, Yeonghojinmi, Danmi, Mogwoo, Boseokheukchal, Jungmo1005, Cheongpungheukchal

Supplementary Table 5. Frequency distribution of rice variety showing types in ligule.

Shape (Ligule)	No. of variety
Partedness	<p>Orae, Hwanggeumchal, Heukgawichal, Geumseong, Joryeongbyeo, Hwanambyeo, Sambaekbyeo, Sangsanbyeo, Junghwabyeo, Namcheonbyeo, Naepungbyeo, Geumobyeo1, Hwasambyeo, Daesanbyeo, Geumobyeo2, Namgangbyeo, Aranghyangchalbyeo, Yonghaebyeo, Sangjuchalbyeo, Gancheokbyeo, Geumnambyeo, Yangjobyeo, Unjangbyeo, Hwasinbyeo, Hyangnambyeo, Samcheonbyeo, Dongan, Nampyeongbyeo, Hwaseonchalbyeo, Dunnaebyeo, Hwajungbyeo, Daeripbyeo1, Nongan, Hyangmibyeo1, Juanbyeo, Daeanybyeo, Dasan, Ansanbyeo, Daejin, Seojin, Hyangmibyeo2, Hwadongbyeo, Geurubyeo, Hwamyongbyeo, Heukjinjubyo, Sura, Gwangan, Anda, Undu, Mihyang, Dongjinchal, Hoan, Inwol, Hwabong, Nongho, Manan, Wonhwang, Sangmi, Jungan, Seolhyangchal, Sobi, Sindongjin, Areum, Sujin, Sampyeong, Taebong, Hwaan, Jinbong, Jeokjinju, Hojin, Manpung, Heukhyang, Junam, Haepyeong, Jungsan, Goami, Jinpum, Saekyehwa, Saesangju, Jongnam, Manwol, Yongan, Seolgaeng, Baekjinju, Seokjeong, Samdeok, Manmi, Geuman, Namil, Taeseong, Daepyeong, Manho, Goami2, Sangok, Saechucheong BIL-3, Dongjin1, Samgwang, Seoeun, Josaengheukchal, Boseokheukchal, Sintoheukmi, Joan, Heukgwang, Hopyeong, Seopyeong, Cheongho, Boseokchal, Hanmaeum, Pungmi, Haepyeongchal, Goun, Gopum, Hwasin1, Sinunbong1, Onnuri, Manna, Dongjin2, Odae1, Juan1, Keunnun, Seon1, Baekjinju1, Gangbaek, Nokyang, Dami, Dasan1, Donghaejinmi, Malgeumi, Sandeuljinmi, Cheongdam, Cheonga, Keunseom, Hangangchal1, Hwanggeumbora, Milkikwin, Noreunjachal, Geumtap, Seonong6, Seonong10, Seonong8, Seonong9, Seonong12, Sanggol, Wonpyeong, Wongwang, Wonmi, Woncheong, Wonpum, Nokwonchal, Wonhae, Jungmo1012, Seonong16, Jinsang, Anmi, Segyejinmi, Manjong, Durujinmi, Daeripjami, Jeokjinjuchal, Suryeojinmi, Dongwon, Jungmo1007, Garak, Chinnong, Jopyeong, Geonganghongmi, Sinbaek Onojjinoijji, Dongbo, Wolbaek, Seolbaek, Migwang, Cheonghaejinmi, Jungmo1001, Geumyeong, Boramchan, Seomyeong, Yoonnongchamssal, Hanareum2, Gangchan, Deurimi2, Syupeojami, Keunnunjami, Jungmo1010, Daechan, Dasan2, Mokyang, Joun, Seonong15, Jinbo, Goami4, Cheongbaekchal, Suan, Seonong14, Jungmo1009, Jungmo1014, Jungmo1011, Jungmo1015, Jungmo1013, Saenuri, Yeonghojinmi, Hiam, Dacheong, Daeyabyeo, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Honggwang, Goami3, Danmi, Mogwoo, Baekokchal, Joami, Jungmo1004, Jungmo1005, Jungmo1006, Jinsumi, Cheongcheongjinmi, Cheongpungheukchal, Pyeongwon, Handeul, Haeoreumi, Hoban, Cheongnam, Jagwangchal, Sinmyeongheukchal, Heuknam, Pungmi1, Nunbora</p>
Sharp	<p>Manchu, Munjang, Hwarang, Pyeongan, Mipum, Heukseonchal, Seongjochal, Hongjinju</p>

Supplementary Table 6. Frequency distribution of flag leaf angle in 243 rice varieties at heading stage.

Flag leaf angle (Heading stage)	No. of variety
Erect	Geumyeong, Manho, Wonpyeong, Joan, Jungmo1011, Taebong, Taeseong, Manna, Manchu, Sangsanbye, Sinunbong1, Ansanbye, Wolbaek, Inwol, Joun, Jungmo1007, Junghwabye, Manan, Munjang, Saesangju, Undu, Wonpum, Jopyeong, Daechan, Jungmo1010, Daeanye, Seomyeong, Seojin, Juanbye, Hojin, Hwarang, Nampyeongbye, Daesanbye, Dongbo, Dongan, Dongjin2, Dongjinchal, Donghaejinmi, Malgeumi, Migwang, Mihyang, Baekjinju, Boramchan, Saekyehwa, Seonong10, Sujin, Woncheong, Junam, Jungmo1013, Cheonga, Cheonghaejinmi, Pyeongan, Hoan, Hwasin1, Gancheokbye, Gangbaek, Gangchan, Goami2, Goami4, Geumnambye, Geumbye, Nongan, Dami, Daejin, Deurimi2, Manpung, Boseokchal, Samgwang, Samdeok, Sanggol, Sangok, Seonong6, Seonong8, Seonong9, Seonong1, Seoeun, Seopyeong, Seolgaeng, Sura, Sinbaek, Anmi, Yangjobye, Yeonghaebye, Wonmi, Wonhwang, Jongnam, Juan1, Jungmo1012, Jungmo1014, Jungmo1009, Jungsan, Cheongho, Hanmaeum, Hanmaeum2, Haepyeong, Haepyeongchal, Hyangnambye, Hwabong, Hwasambye, Hwaan, Hwajungbye, Geonganghongmi, Goami, Geuman, Geumbye2, Namgangbye, Nokyang, Daeripbye1, Daepyeong, Manmi, Mokyang, Sampyeong, Saechucheong BIL-3, Seonong15, Suan, Sindongjin, Arangyangchalbye, Onojinoijji, Wonchu, Jungan, Jinbo, Jinsang, Gwangan, Noreunjachal, Dongjin1, Manjong, Seonong14, Sobi, Yeongan, Jinpum, Keunnun, Hopyeong, Heukgwang, Nongho, Mipum, Syupeojami, Suryeojinmi, Heukhyang, Baekjinju1, Jungmo1015, Saenuri, Hiami, Dacheong, Daeyabye, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Honggwang, Goami3, Baekokchal, Joami, Jungmo1004, Jungmo1006, Jinsumi, Cheongcheongjinmi, Pyeongwon, Handeul, Haoreumi, Hoban, Cheongnam, Gopum, Sandeuljinmi, Samcheonbye, Jeokjinju, Jinbong, Hwanggeumbora, Heukseonchal, Geumseong, Sambaekbye, Sangjuchalbye, Wonhae, Hyangmibye2, Sangmi, Jeokjinjuchal, Heukjinjuby, Jagwangchal, Hwanggeumchal, Hwamyongbye, Naepungbye, Dasan, Dasan1, Seonong16, Segyejinmi, Chinnong, Keunseom, Hyangmibye1, Namcheonbye, Nokwonchal, Dasan2, Seonong12, Areum, Anda, Orae, Yoonnongchamssal, Jungmo1001, Cheongdam, Pungmi, Hangangchal1, Hongjinju, Hwanambye, Hwadongbye, Hwaseonchalbye, Seongjochal, Geumtap, Nunbora, Manwol, Yeonghojinmi, Danmi, Mogwoo, Jungmo1005, Cheongpungheukchal, Heukgawichal, Sinmyeongheukchal, Josaengheukchal, Boseokheukchal, Heuknam, Sintoheukmi
Semierect	Pungmi1, Durujinmi, Unjangbye, Seolbaek, Goun, Odae1, Geurubye, Namil, Milkikwin, Seolhyangchal, Dongwon, Keunnunjami, Seokjeong, Onnuri, Wongwang, Hwasinbye, Garak, Daeripjami, Dunnaebye, Cheongbaekchal, Joryeongbye
Horizontal	0

Supplementary Table 7. Frequency distribution of flag leaf angle in 243 rice varieties at yellow ripe stage.

Flag leaf angle (Yellow ripe stage)	No. of variety
Erect	Pungmi1, Goun, Wongwang, Dongwon, Daeripjami, Gopum, Jungmo1011, Taeseong, Manna, Manchu, Sangsanbyeol, Ansanbyeol, Wolbaek, Inwol, Joun, Jungmo1007, Junghwabyeol, Manan, Munjang, Saesangju, Daechan, Jungmo1010, Daeanbyeol, Seomyeong, Juanbyeol, Hojin, Hwarang, Daesanbyeol, Dongbo, Dongan, Dongjin2, Dongjinchal, Donghaejinmi, Malgeumi, Migwang, Mihyang, Baekjinju, Boramchan, Saekyehwa, Seonong10, Sujin, Woncheong, Junam, Jungmo1013, Cheonga, Cheonghaejinmi, Pyeongan, Hoan, Hwasin1, Gancheokbyeol, Gangbaek, Gangchan, Goami2, Goami4, Geumnambyeol, Geumbyeol1, Nongan, Dami, Daejin, Deurimi2, Manpung, Boseokchal, Samgwang, Samdeok, Sanggol, Sangok, Seonong6, Seonong8, Seonong9, Seonan1, Seopyeong, Seolgaeng, Sura, Sinbaek, Anmi, Yangjobyeol, Wonmi, Wonhwang, Jongnam, Juan1, Jungmo1012, Jungmo1014, Jungmo1009, Jungsan, Hanmaeum, Hanareum2, Hyangnambyeol, Hwabong, Hwaan, Hwajungbyeol, Geonganghongmi, Goami, Geuman, Geumbyeol2, Nokyang, Manmi, Mokyang, Saechucheong BIL-3, Suan, Sindongjin, Arangyangchalbyeol, Onojjinoijji, Wonchu, Jungan, Jinsang, Gwangan, Noreunjachal, Sobi, Yeongan, Jinpum, Keunnun, Hopyeong, Heukgwang, Nongho, Mipum, Syupeojami, Suryeojinmi, Heukhyang, Baekjinju1, Jungmo1015, Sandeuljinmi, Samcheonbyeol, Hwanggeumbora, Geumseong, Sambaekbyeol, Sangjuchalbyeol, Hyangmibyeol2, Sangmi, Jeokjinjuchal, Hwanggeumchal, Hwamyongbyeol, Naepungbyeol, Dasan, Dasan1, Seonong16, Segyejinmi, Chinnong, Keunseom, Hyangmi1, Namcheonbyeol, Nokwonchal, Dasan2, Seonong12, Areum, Anda, Orae, Yoonnongchamssal, Jungmo1001, Cheongdam, Hangangchal1, Hongjinju, Hwanambyeol, Hwadongbyeol, Hwaseonchalbyeol, Seongjochal, Geumtap, Nunbora, Saenuri, Hiam, Dacheong, Daeyabyeol, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodel, Goami3, Baekokchal, Jungmo1004, Jungmo1006, Jinsumi, Cheongcheongjinmi, Pyeongwon, Cheongnam, Yeonghojinmi, Danmi, Mogwoo, Jungmo1005, Cheongpungheukchal, Seoeun, Jopyeong, Sinmyeongheukchal, Heukgawichal, Josaengheukchal, Boseokheukchal, Heuknam, Sintoheukmi
Semierect	Durujinmi, Unjangbyeol, Geurubyeol, Seolbaek, Namil, Seokjeong, Onnuri, Hwasinbyeol, Milkikwin, Seolhyangchal, Garak, Dunnaebyeol, Joryeongbyeol, Geumyeong, Manho, Wonpyeong, Joan, Taebong, Sinunbong1, Wonpum, Seojin, Nampyeongbyeol, Yeonghaebyeol, Cheongho, Haepyeong, Haepyeongchal, Hwasambyeol, Namgangbyeol, Daepyeong, Sampyeong, Seonong15, Jinbo, Dongjin1, Manjong, Seonong14, Jeokjinju, Jinbong, Heukseonchal, Wonhae, Jagwangchal, Pungmi, Manwol, Joami, Handeul, Haeoreumi
Horizontal	Odae1, Keunnunjami, Cheongbaekchal, Honggwang, Undu, Daeripbyeol1, Heukjinju, Hoban

Supplementary Table 8. Frequency distribution of culm thickness in 243 rice varieties.

Thickness (Culm)	No. of variety
Thin	Unjangbyeo, Geurubyeo, Milkikwin, Dunnaebyeo, Geumyeong, Wolbaek, Manpung, Seolbaek, Joami, Onnuri, Manwol, Heukjinjubyoo, Joun, Inwol, Taeseong, Juanbyeoo, Yongan, Jongnam, Sandeuljinmi, Samcheonbyeoo, Sambaekbyeoo, Unmi, Seoun, Honggwang
Medium	Durujinmi, Namil, Seokjeong, Hwasinbyeoo, Seolhyangchal, Garak, Joryeongbyeoo, Manho, Wonpyeong, Joan, Taebong, Sinunbong1, Wompum, Seojin, Nampyeongbyeoo, Cheongho, Haepyeong, Hwasambyeoo, Namgangbyeoo, Daepyeong, Sampyeong, Jinbo, Dongjin1, Manjong, Seonong14, Jeokjinju, Jinbong, Heukseonchal, Wonhae, Pungmi, Handeul, Undu, Hoban, Goun, Wongwang, Dongwon, Gopum, Jungmo1011, Manchu, Sangsanbyeoo, Ansanbyeoo, Junghwabyeo, Manan, Munjang, Saesangju, Daechan, Jungmo1010, Daeanbyeoo, Seomyeong, Hojin, Hwarang, Daesanbyeoo, Dongbo, Dongan, Donghaejinmi, Malgeumi, Migwang, Mihyang, Baekjinju, Boramchan, Saekyehwa, Seonong10, Woncheong, Junam, Jungmo1013, Cheonga, Hoan, Hwasin1, Gancheokbyeoo, Gangbaek, Gangchan, Goami2, Geumnambyeoo, Geumbyeoo1, Nongan, Daejin, Deurimi2, Boseokchal, Samgwang, Samdeok, Sanggol, Sangok, Seonong6, Seonong8, Seonong9, Seoan1, Seopyeong, Seolgaeng, Sinbaek, Anmi, Yangjobyeoo, Wonmi, Juan1, Jungmo1012, Jungmo1014, Jungsan, Hanmaeum, Hyangnambyeoo, Hwabong, Hwaan, Hwajungbyeoo, Geonganghongmi, Goami, Geuman, Geumbyeoo2, Manmi, Saechucheong BIL-3, Suan, Aranghyangchalbyeoo, Wonchu, Jungan, Jinsang, Gwangan, Noreunjachal, Sobi, Jinpum, Hopyeong, Heukgwang, Mipum, Heukhyang, Baekjinju1, Jungmo1015, Hwanggeumbora, Geumseong, Sangjuchalbyeoo, Sangmi, Jeokjinjuchal, Naepungbyeoo, Seonong16, Keunseom, Hyangmibyeoo1, Namcheonbyeoo, Nokwonchal, Areum, Cheongdam, Hongjinju, Hwadongbyeoo, Hwaseonchalbyeoo, Saenuri, Hiami, Dacheong, Daeyabyeo, Deuraechan, Boseok, Seogan, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Goami3, Baekokchal, Jungmo1004, Jinsumi, Cheongcheongjinmi, Pyeongwon, Cheongnam, Yeonghojinmi, Danmi, Noku, Jungmo1005, Jopyeong, Heukgawichal, Josaengheukchal, Sintoheukmi
Thick	Yeonghaebyeoo, Haepyeongchal, Seonong15, Jagwangchal, Haeoreumi, Odae1, Keunnunjami, Cheongbaekchal, Daeripbyeoo1, Pungmi1, Daeripjami, Manna, Jungmo1007, Dongjin2, Dongjinchal, Sujin, Cheonghaejinmi, Pyeongan, Goami4, Dami, Sura, Wonhwang, Jungmo1009, Hanareum2, Nokyang, Nokyang, Sindongjin, Onojjinoijji, Keunnun, Nongho, Syupeojami, Suryeojinmi, Hyangmibyeoo2, Hwanggeumchal, Hwamyongbyeoo, Dasan, Dasan1, Segyejinmi, Chinnong, Dasan2, Seonong12, Anda, Orae, Yoonnongchamssal, Jungmo1001, Hangangchal1, Hwanambyeoo, Seongjochal, Geumtap, Nunbora, Jungmo1006, Cheongpungheukchal, Sinmyeongheukchal, Boseokheukchal, Heuknam

Supplementary Table 9. Frequency distribution of culm attitude at maturity in 243 rice varieties.

Attitude (Culm)	No. of variety
Erect	Goun, Wongwang, Dongwon, Daeripjami, Gopum, Jungmo1011, Taeseong, Manna, Ansanbyeo, Inwol, Joun, Junghwabyeo, Manan, Saesangju, Daechan, Jungmo1010, Daeanbyeo, Seomyeong, Juanbyeo, Hojin, Hwarang, Daesanbyeo, Dongbo, Dongan, Dongjin2, Dongjinchal, Donghaejinmi, Malgeumi, Migwang, Mihyang, Baekjinju, Boramchan, Saekyehwa, Seonong10, Sujin, Woncheong, Junam, Jungmo1013, Cheonga, Cheonghaejinmi, Pyeongan, Hoan, Gancheokbyeo, Gangbaek, Gangchan, Goami2, Goami4, Geumnambyeo, Geumobyeo1, Nongan, Dami, Daejin, Deurimi2, Boseokchal, Samgwang, Samdeok, Sangok, Seonong6, Seonong8, Seonong9, Seonan1, Seopyeong, Sura, Anmi, Yangjobyeo, Wonmi, Wonhwang, Jongnam, Juan1, Jungmo1012, Jungmo1014, Jungmo1009, Jungsan, Hanmaeum, Hanareum2, Hyangnambyeo, Hwabong, Hwaan, Hwajungbyeo, Geonganghongmi, Goami, Geuman, Geumobyeo2, Nokyang, Manmi, Mokyang, Saechucheong BIL-3, Suan, Sindongjin, Aranghyangchalbyeo, Onojjinoijji, Wonchu, Jungan, Jinsang, Gwangan, Noreunjachal, Sobi, Yeongan, Jinpum, Keunnun, Hopyeong, Heukgwang, Nongho, Mipum, Syupeojami, Suryeojinmi, Heukhyang, Baekjinju1, Jungmo1015, Sandeuljinmi, Samcheonbyeo, Hwanggeumbora, Geumseong, Sambaekbyeo, Hyangmibyeo2, Sangmi, Jeokjinjuchal, Hwanggeumchal, Hwamyongbyeo, Naepungbyeo, Dasan, Dasan1, Seonong16, Segyejinmi, Chinnong, Hyangmi1, Namcheonbyeo, Nokwonchal, Dasan2, Areum, Anda, Orae, Yoonnongchamssal, Jungmo1001, Cheongdam, Hongjinju, Hwanambyeo, Hwadongbyeo, Hwaseonchalbyeo, Seongjochal, Geumtap, Saenuri, Hiami, Dacheong, Daeyabyeo, Deuraechan, Boseok, Seogan, Unmi, Jungmo1008, Jinbaek, Honong, Hwanggeumnodeul, Goami3, Baekokchal, Jungmo1004, Jungmo1006, Jinsumi, Cheongcheongjinmi, Yeonghojinmi, Danmi, Mogwoo, Jungmo1005, Cheongpungheukchal, Seoeun, Jopyeong, Heukgawichal, Josaengheukchal, Heuknam, Sintoheukmi, Onnuri, Hwasinbyeo, Seolhyangchal, Garak, Joryeongbyeo, Wonpyeong, Sinunbong1, Wonpum, Seojin, Nampyeongbyeo, Yeonghaebyeo, Cheongho, Hwasambyeo, Namgangbyeo, Daepyeong, Sampyeong, Seonong15, Jinbo, Dongjin1, Manjong, Seonong14, Jeokjinju, Jinbong, Heukseonchal, Wonhae, Jagwangchal, Pungmi, Manwol, Handeul, Haeoreumi, Odae1, Keunnunjami, Honggwang, Undu, Daeripbyeol, Heukjinju
Semierect	Cheongbaekchal, Pungmi1, Jungmo1007, Hangangchal1, Nunbora, Sinmyeongheukchal, Boseokheukchal, Durujinmi, Seokjeong, Joan, Taebong, Haepyeong, Heukseonchal, Manchu, Sangsanbyeo, Munjang, Hwasin1, Sanggol, Seolgaeng, Sinbaek, Sangjuchalbyeo, Keunseom, Unjangbyeo, Geurubyeo, Milkikwin, Dunnaebyeo, Geumyeong, Wolbaek, Manpung
Medium	Seolbaek, Namil, Manho, Joami, Hoban, Seonong12, Pyeongwon, Cheongnam

Supplementary Figure 1. Continued.

Variety Name		Phenotype	Panicle	Brown rice
Nampyeongbyeo	남평벼
Neongeongbyeo	넴경벼
Nokwongchal	녹원찰

Nokyang	녹양
Nongan	농안
Nongho	농호

Noreunjachal	노른자찰
Nunbora	눈보라
Odae1	오대1호

Onnuri	온누리
Onojjinoijji	오노찌노이찌
Orae	오래

Pungmi	풍미
Pungmi1	풍미1호
Pyeongang	평안

Pycongwon	평원
SaechucheongBIL-3	새추청 BIL-3
Saegye-hwa	새계화

Saenuri	새누리
Saesangju	새상주
Sambaekbyeon	삼백벼

Samcheonbyeo	삼천벼
Samgwang	삼광
Samyeong	삼평

Sandeok	삼덕
Sandeuljinmi	산들진미
Sanggal	상골

Sangjuchalbyeo	상주찰벼
Sangmi	상미
Sangok	상옥

Sangsanbyeo	상산벼
Segyejinmi	세계진미
Seoan1	서안1호

Seoeun	서은
Seogan	서간
Seojin	서진

Seokjeong	석정
Seolbaek	설백
Seolgaeng	설강

Seolhyangchal	설향찰
Seomyeong	서명
Seongjochal	성조찰

Seonong10	서농10호
Seonong12	서농12호
Seonong14	서농14

Seonong15	서농15
Seonong16	서농16
Seonong6	서농6호

Seonong8	서농8호
Seonong9	서농9호
Seopyeong	서평

Sinbaek	신백
Sindongjin	신동진
Sinmyeongheukcha 1	신명흑찰

Sintoheukmi	신토흑미
Sinunbong1	신운봉1호
Sobi	소비

Suan	수안
Sujin	수진
Sura	수라

Suryeojinmi	수려진미
Syupeojami	슈퍼자미
Taebong	태봉

Taeseong	태성
Undu	운두
Unjangbyeon	운장벼

Unmi	운미
Wolbaek	월백
Woncheong	원청

Wonchu	원추
Wongwang	원광
Wonhae	원해

Wonhwang	원황
Wonmi	원미
Wonpum	원품

Wonpyeong	원평
Yangjobyeo	양조벼
Yeongan	영안

Yeonghaebyeo	영해벼
Yeonghojinmi	영호진미
Yoonnongchamssal	윤농참쌀